

KMS-weights on C^* -algebras

J. Kustermans ¹

Institut for Matematik og Datalogi

Odense Universitet

Campusvej 55

5230 Odense M

Denmark

e-mail : johank@imada.ou.dk

April 1997

Abstract

In this paper, we build a solid framework for the use of KMS-weights on C^* -algebras. We will use another definition than the one introduced by Combes in [4], but we will prove that they are equivalent. However, the subject of KMS-weights is approached from a somewhat different angle. We introduce a construction procedure for KMS-weights, prove the most important properties of them, construct the tensor product of KMS-weights and construct weights which are absolutely continuous to a given weight.

Introduction

In [3], Combes studied lower semi-continuous weights on C^* -algebras and proved the major result that a lower semi-continuous weight can be approximated by the positive linear functionals which are majorated by it.

At the moment, these lower semi-continuous weights seem not to behave well enough to be used in an efficient way. So it is natural to search for extra conditions on lower semi-continuous weights which makes them manageable. An attempt in this direction was undertaken by Jan Verding in [22] where he introduced the so-called regular weights. These are lower semi-continuous weights which have a well-behaved truncating net.

It turns out that this regularity condition is also very useful in the theory of C^* -valued weights (see [10]).

There is however a class of lower semi-continuous weights which behave very well and were introduced by Combes in definition 4.1 of [4], the so-called KMS-weights:

Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A such that there exists a norm continuous one-parameter group σ on A satisfying:

1. We have for every $t \in \mathbb{R}$ that $\varphi\sigma_t = \varphi$.
2. For every $x, y \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$, there exists a bounded continuous function f from $S(i)$ into $\mathbb{C}$ which is analytic on $S(i)^0$ and such that:
 - We have for every $t \in \mathbb{R}$ that $f(t) = \varphi(\sigma_t(x)y)$.
 - We have for every $t \in \mathbb{R}$ that $f(t+i) = \varphi(y\sigma_t(x))$.

¹Research Assistant of the National Fund for Scientific Research (Belgium)

Then φ is called a KMS-weight on A , σ is called a modular group for φ .

Here, $S(i)$ denotes the horizontal strip in the complex plane between 0 and i .

By a proof of Combes in [5], we know that such a KMS-weight can be extended to normal KMS-weight in the GNS-representation of φ . In this way, we can make use of the very rich theory of normal weights on von Neumann algebras and use for instance the Radon Nikodym theorem for normal weights (see [14]).

In short, we get a class of weights which posses the same rich structure as the Borel measures on locally compact spaces.

The last years, there is a lot of interest for lower-semicontinuous weights in general (and KMS-weights in particular) from people working in the field of C^* -algebraic quantum groups. This because of the role of the left Haar weight in C^* -algebraic quantum group theory.

At the moment, it is not yet clear whether the left Haar weight should be a KMS-weight. There are however some indications that this will be the case :

- Masuda, Nakagami & Woronowicz are momentarily working on a definition for reduced C^* -algebraic quantum group and their left Haar weight will be a KMS-weight.
- In all the known examples, the left Haar weight turns out to satisfy the KMS-condition.
- A. Van Daele proves in [21] that the left Haar functional on an algebraic quantum group satisfies automatically some sort of weak KMS-condition.
- It is proven in [8] that this weak KMS-condition implies that the left Haar weight on the reduced C^* -algebraic group arising from the algebraic one, is a KMS-weight.
- In [11], we prove that this KMS-condition on this reduced C^* -algebraic quantum group implies that the left Haar weight on the universal C^* -algebraic quantum group is also KMS. It appears that the same principle can be used to prove such a result in the general cas.

Especially the universal case necessitates to investigate KMS-weights in a C^* -algebraic framework.

In this paper, we will work with another definition of a KMS-weight than the one mentioned above but we will show that both are equivalent. One of the reasons to use the other definition is the fact that Masuda, Nakagami & Woronowicz seemed to be going to use this other definition in their approach to quantum groups.

The main aim of this paper is to build a solid framework for KMS-weights on C^* -algebras. We believe that it gives a technically useful overview of KMS-weights in the C^* -algebra picture.

Most of the results will look familiar to the ones known in normal weight theory but will be proven within the C^* -algebra framework.

In the first section, we fix notations and give an overview of the results about one-parameter representations and their analytic continuations.

We give the definition of a KMS-weight in the second section. At the same time, we prove some useful properties concerning lower semi-continuous weights.

In section 3, we have gathered some results from [22] about the construction of a weight starting from a GNS-construction.

The fourth section concerns the behaviour of one-parameter representations which are relatively invariant with respect to some closed mapping.

In the fifth section, we prove that a closed linear mapping from within a C^* -algebra into a Hilbert space and satisfying certain KMS-characteristics gives rise to a KMS-weight on this C^* -algebra.

The most important technical properties concerning KMS-weights are collected in the sixth section.

In section 7, we construct the tensor product of KMS-weights.

Given a KMS-weight φ and a strictly positive element δ affiliated to the C^* -algebra which is relatively invariant under a modular group of φ , we construct the weight $\varphi(\delta^{\frac{1}{2}} \cdot \delta^{\frac{1}{2}})$ in section 8. We have to use another method than the one used by Pedersen & Takesaki because of the relative invariance.

Let us now fix some terminology and conventions.

Consider a complex number z , then $S(z)$ will denote the horizontal strip $\{c \in \mathbb{C} \mid \text{Im } c \in [0, \text{Im } z]\}$. The interior of $S(z)$ will be denoted by $S(z)^0$.

The domain of a mapping f will be denoted by $D(f)$, its image by $\text{Ran } f$.

For any Banach space E , we denote the set of bounded operators on E by $B(E)$ and the topological dual by E^* .

Let A be a C^* -algebra. We denote the multiplier algebra of A by $M(A)$.

Any element $\omega \in A^*$ has a unique strictly continuous linear extension to $M(A)$ and we denote this extension by $\overline{\omega}$. We put $\omega(x) = \overline{\omega}(x)$ for every $x \in M(A)$.

For a good introduction into elements affiliated with A , we refer to [12] and [24].

Whenever we speak of a positive element affiliated element with A , we mean a (possibly unbounded) positive and selfadjoint element affiliated with A . The same remark applies to positive operators in Hilbert spaces. A positive element affiliated to A is called strictly positive if it has dense range. As usual, we can raise a strictly positive affiliated element to any complex power.

Consider a Hilbert space H . Let π a non-degenerate $*$ -homomorphism from A into $B(H)$. Then π has a unique extension to a $*$ -representation $\overline{\pi}$. We put $\pi(x) = \overline{\pi}(x)$ for every $x \in M(A)$.

If T is an element affiliated with A , then $\pi(T)$ denotes the closed densely defined operator in H such that $\pi(D(T))$ is a core for $\pi(T)$ and $\pi(T)(\pi(a)v) = \pi(T(a))v$ for every $a \in D(T)$ and $v \in H$.

In the following, $\circ$ will denote the composition of mappings and elements of $M(A)$ will be considered as linear mappings from A into A .

Consider an element T affiliated with A and $a \in M(A)$. Then

- We call a a left multiplier of T if $a \circ T$ is bounded as a mapping from A into A and $\overline{a \circ T}$ belongs to $M(A)$. In this case, we put $aT = \overline{a \circ T}$
- We call a a right multiplier of T if $D(T \circ a) = A$. You can prove in this case that $T \circ a$ belongs to $M(A)$ and we put $Ta = T \circ a$.

It is not very difficult to prove that a is a left multiplier of T if and only if a is a right multiplier of T^* . If this is the case, we have that $(aT)^* = T^*a^*$.

We will freely use calculation rules involving this left and right multipliers, e.g. Let $a, x \in M(A)$ and T an element affiliated with A . If a is a left multiplier of T , then xa is a left multiplier of T and $(xa)T = x(aT)$.

Consider elements S, T affiliated with A and $a \in M(A)$. Then we call a a middle multiplier of S, T if $D(S \circ a \circ T) = D(T)$, $S \circ a \circ T$ is bounded and $\overline{S \circ a \circ T}$ belongs to $M(A)$. In this case, we put $SaT = \overline{S \circ a \circ T}$.

1 One-parameter representations

In this first section, we recall the definition of one-parameter representations and an overview of the most important results. Most of the proofs can be found in [9]. A standard reference for one-parameter representation is [25].

Whenever we use the notion of integrability of a function with values in a Banach space, we mean the strong form of integrability (e.g. Analysis II, S. Lang) :

Consider a measure space $(X, \mathcal{M}, \mu)$ and a Banach space E .

- It is obvious how to define integrability for step functions from X into E .
- Let f be a function from X into E . Then f is μ -integrable if and only if there exists a sequence of integrable step functions $(f_n)_{n=1}^\infty$ from X into E such that :
 1. We have for almost every $x \in X$ that $(f_n(x))_{n=1}^\infty$ converges to $f(x)$ in the norm topology.
 2. The sequence $(f_n)_{n=1}^\infty$ is convergent in the L_1 -norm.

In this case, the sequence $(\int f_n d\mu)_{n=1}^\infty$ is convergent and the integral of $\int f d\mu$ is defined to be the limit of this sequence (Of course, one has to prove that this limit is independent of the choice of the sequence $(f_n)_{n=1}^\infty$).

- It is possible to define a form of μ -measurability (see Lang) in such a way that a function f is integrable if and only if the function f is measurable and $\|f\|$ is integrable.

We start this section with a basic but very useful lemma.

Lemma 1.1 *Consider Banach spaces E, F , Λ a closed linear mapping from within E into F . Let f be a function from $\mathbb{R}$ into $D(\Lambda)$ such that*

- *f is integrable.*
- *The function $\mathbb{R} \rightarrow F : t \mapsto \Lambda(f(t))$ is integrable.*

Then $\int f(t) dt$ belongs to $D(\Lambda)$ and $\Lambda(\int f(t) dt) = \int \Lambda(f(t)) dt$.

Proof : Define G as the graph of the mapping Λ . By assumption, we have that G is a closed subspace of $E \oplus F$. Next, we define the mapping g from $\mathbb{R}$ into G such that $g(t) = (f(t), \Lambda(f(t)))$ for every $t \in \mathbb{R}$. It follows that g is integrable and

$$\int g(t) dt = \left(\int f(t) dt, \int \Lambda(f(t)) dt \right).$$

Because G is a closed subspace of $E \oplus F$, we have that $\int g(t) dt$ belongs to G . This implies that $(\int f(t) dt, \int \Lambda(f(t)) dt)$ belongs to G . ■

When we speak of analyticity, we will always mean norm analyticity. But we have the following well known result (which follows from the Uniform Boundedness principle).

Result 1.2 *Consider a Banach space E and a subspace F of E^* such that $\|x\| = \sup\{\|\omega(x)\| \mid \omega \in F \text{ with } \|\omega\| \leq 1\}$ for every $x \in E$. Let O be an open subset of $\mathbb{C}$ and f a function from O into E . Then f is analytic $\Leftrightarrow$ We have for every $\omega \in F$ that $\omega \circ f$ is analytic.*

Notice that the result is true for $F = E^*$.

1.1 One-parameter representations on Banach spaces

We introduce the notion of strongly continuous one-parameter groups and discuss the analytic continuations of them. The standard reference for the material in this section is [25].

Terminology 1.3 *Consider a Banach space E . By a one-parameter representation on E , we will always mean a mapping α from $\mathbb{R}$ into $B(E)$ such that :*

- *We have for every $s, t \in \mathbb{R}$ that $\alpha_{s+t} = \alpha_s \alpha_t$.*
- *$\alpha_0 = \iota$*
- *We have for every $t \in \mathbb{R}$ that $\|\alpha_t\| \leq 1$.*

We call α strongly continuous $\Leftrightarrow$ We have for every $a \in E$ that the mapping $\mathbb{R} \rightarrow E : t \mapsto \alpha_t(a)$ is continuous.

This definition implies for every $t \in \mathbb{R}$ that α_t is invertible in $B(E)$, that $(\alpha_t)^{-1} = \alpha_{-t}$ and that $\|\alpha_t\| = 1$.

Remark 1.4 We would like to mention the following special cases :

- If H is a Hilbert space and u is a strongly continuous one-parameter representation on H such that u_t is unitary for every $t \in \mathbb{R}$, we call u a strongly continuous unitary one-parameter group on H .
- If A is a C^* -algebra and α is a strongly continuous one-parameter representation on A such that α_t is a C^* -automorphism on A for every $t \in \mathbb{R}$, we call α a norm continuous one-parameter group on A .
- If A is a C^* -algebra and u is a strongly continuous one-parameter representation on A such that u_t is a unitary element in $M(A)$ for every $t \in \mathbb{R}$, we call u a strictly continuous unitary one-parameter group on A .

Now we will summarize the theory of analytic continuations of such norm continuous one-parameter representations. For the most part of this subsection, we will fix a Banach space E and a strongly continuous one-parameter representation α on E .

Definition 1.5 *Consider $z \in \mathbb{C}$. We define the mapping α_z from within E into E such that :*

- *The domain of α_z is by definition the set $\{a \in E \mid \text{There exists a function } f \text{ from } S(z) \text{ into } E \text{ such that}$*
 - 1. f is continuous on $S(z)$*
 - 2. f is analytic on $S(z)^0$*
 - 3. We have that $\alpha_t(a) = f(t)$ for every $t \in \mathbb{R}$ }.*
- *Choose a in the domain of α_z and f the function from $S(z)$ into E such that*
 - 1. f is continuous on $S(z)$*
 - 2. f is analytic on $S(z)^0$*
 - 3. We have that $\alpha_t(a) = f(t)$ for every $t \in \mathbb{R}$*

Then we have by definition that $\alpha_z(a) = f(z)$.

In the case that z belongs to $\mathbb{R}$, this definition of α_z corresponds with α_z which we started from. Therefore, the notation is justified.

Remark 1.6 • Consider $z \in \mathbb{C}$ and $y \in S(z)$. Then it is clear that $D(\alpha_z)$ is a subset of $D(\alpha_y)$.

- It is also clear that $D(\alpha_y) = D(\alpha_z)$ for $y, z \in \mathbb{C}$ with $\text{Im } y = \text{Im } z$.
- Let $z \in \mathbb{C}$ and $a \in D(\alpha_z)$. Then the function $S(z) \rightarrow E : u \mapsto \alpha_u(a)$ is continuous on $S(z)$ and analytic on $S(z)^0$ (because this function must be equal to the function f from the definition).
- Consider an element a in E . We say that a is analytic with respect to α if a belongs to $D(\alpha_z)$ for every $z \in \mathbb{C}$. If a is analytic with respect to α , then the function $\mathbb{C} \rightarrow E : u \mapsto \alpha_u(a)$ is analytic.

Now we give a list of the most important properties of these analytic continuations. The most important tool for proving the results is the Phragmen-Lindelof theorem.

Proposition 1.7 *The mapping α_z is a closed linear operator in E which is densely defined and has dense range.*

We would like to mention that the proof of the closedness is substantially easier than in the case of strongly continuous one-parameter groups on von Neumann algebras (see [25]).

Result 1.8 *Consider $z \in \mathbb{C}$ and $a \in D(\alpha_z)$. Then the function $S(z) \rightarrow E : u \mapsto \alpha_u(a)$ is bounded.*

Proposition 1.9 *Consider $z \in \mathbb{C}$ and $t \in \mathbb{R}$. Then $\alpha_z \alpha_t = \alpha_t \alpha_z = \alpha_{z+t}$. Furthermore, the following equalities hold :*

- We have that $\alpha_t(D(\alpha_z)) = D(\alpha_z)$ and $\alpha_t(\text{Ran } \alpha_z) = \text{Ran } \alpha_z$.
- We have that $D(\alpha_{z+t}) = D(\alpha_z)$ and $\text{Ran } \alpha_{z+t} = \text{Ran } \alpha_z$.

Proposition 1.10 *Consider $z \in \mathbb{C}$. Then α_z is injective and $(\alpha_z)^{-1} = \alpha_{-z}$. Therefore $\text{Ran } \alpha_z = D(\alpha_{-z})$ and $D(\alpha_z) = \text{Ran } \alpha_{-z}$.*

Proposition 1.11 *Consider $y, z \in \mathbb{C}$. Then*

1. $\alpha_y \alpha_z \subseteq \alpha_{y+z}$.
2. If y and z lie at the same side of the real axis, we have that $\alpha_y \alpha_z = \alpha_{y+z}$.

As usual, we smear elements to construct elements which behave well with respect to α .

Notation 1.12 *Consider $a \in E$, $n > 0$ and $z \in \mathbb{C}$. Then we define the element $a(n, z)$ in E such that*

$$a(n, z) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t - z)^2) \alpha_t(a) dt .$$

Then we have that $\|a(n, z)\| \leq \|a\| \exp(n^2(\text{Im } z)^2)$.

For $n > 0$ and $z \in \mathbb{C}$, we will use the notation

$$a(n) = a(n, 0) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) dt .$$

Proposition 1.13 *Let $a \in E$, $n > 0$ and $z \in \mathbb{C}$. Then $a(n, z)$ is analytic with respect to α and $\alpha_y(a(n, z)) = a(n, z + y)$ for every $y \in \mathbb{C}$.*

Proposition 1.14 *Consider $n > 0$, $y, z \in \mathbb{C}$ and $a \in D(\alpha_y)$. Then $\alpha_y(a(n, z)) = \alpha_y(a)(n, z)$.*

A proof of the next result can be found in [23].

Proposition 1.15 *Let z be a complex number. Consider a dense subset K of E . Then the set $\langle a(n) \mid a \in K, n \in \mathbb{N} \rangle$ is a core for α_z .*

We end this subsection with some special cases :

The first one is a familiar one and the proof can be found in corollary 9.21 of [18].

Proposition 1.16 *Consider a Hilbert space H and an injective positive operator M in H . Define the strongly continuous unitary one-parameter group u on H such that $u_t = M^{it}$ for every $t \in \mathbb{R}$. Then $u_z = M^{iz}$ for every $z \in \mathbb{C}$.*

The next result is a C^* -version of the previous one.

Proposition 1.17 *Consider a C^* -algebra A and a strictly positive element δ affiliated with A . Define the strictly continuous unitary one-parameter group u on A such that we have for every $t \in \mathbb{R}$ that $u_t = \delta^{it}$ considered as a left multiplier. Then $u_z = \delta^{iz}$ for every $z \in \mathbb{C}$.*

The previous result implies immediately the following result.

Proposition 1.18 *Consider a C^* -algebra A and a strictly positive element δ affiliated with A . Define the strictly continuous unitary one-parameter group u on A such that we have for every $t \in \mathbb{R}$ that $u_t = \delta^{it}$ considered as a right multiplier. Let z be a complex number and a an element in A , then :*

- *The element a belongs to $D(u_z) \Leftrightarrow a$ is a left multiplier of δ^{iz} and $a\delta^z$ belongs to A .*
- *If a belongs to $D(u_z)$, then $u_z(a) = a\delta^{iz}$.*

The following proposition is also a C^* -version of a known result in Hilbert space theory (see proposition 9.24 of [18]).

Proposition 1.19 *Consider a C^* -algebra A and a strictly positive element δ affiliated with A . Define the norm continuous one-parameter group α on A such that $\alpha_t(a) = \delta^{-it}a\delta^{it}$ for every $t \in \mathbb{R}$ and $a \in A$. Let z be a complex number and a an element in A , then :*

- *The element a belongs to $D(\alpha_z) \Leftrightarrow a$ is a middle multiplier of δ^{-iz} , δ^{iz} and $\delta^{-iz}a\delta^{iz}$ belongs to A .*
- *If a belongs to $D(\alpha_z)$, then $\alpha_z(a) = \delta^{-iz}a\delta^{iz}$.*

A proof of the last 3 propositions can be found in [9].

1.2 One-parameter groups on C^* -algebras

For this section, we will fix a norm-continuous one-parameter group α on a C^* -algebra A . We will investigate the properties of the analytic continuations in this case a little bit further. Proofs of the results in this section can be found in [9].

By the remarks of the previous section, we have for any $z \in \mathbb{C}$ a closed linear operator α_z from within A into A which is generally unbounded.

Because α_t is a $*$ -homomorphism for every $t \in \mathbb{R}$, we have the two following algebraic properties.

Proposition 1.20 *The mapping α_z is a multiplicative linear operator in A .*

Proposition 1.21 *Consider $z \in \mathbb{C}$. Then $D(\alpha_{\bar{z}}) = D(\alpha_z)^*$ and $\text{Ran } \alpha_{\bar{z}} = (\text{Ran } \alpha_z)^*$. We have for every $a \in D(\alpha_z)$ that $\alpha_z(a)^* = \alpha_{\bar{z}}(a^*)$.*

This implies for all $z \in \mathbb{R}i$ and $a \in D(\alpha_z)$ that $\alpha_z(a)^*$ belongs to $D(\alpha_z)$ and $\alpha_z(\alpha_z(a)^*)^* = a$.

For the rest of this section, we want to concentrate on strictly analytic continuations of α .

Concerning analyticity, we have the following result. The proof of this fact uses result 1.2 and the fact that every continuous linear functional on A is of the form $a\omega$ with $\omega \in A^*$ and $a \in A$.

Result 1.22 *Consider an open subset O of the complex plane, f a function from O into $M(A)$. Then f is analytic $\Leftrightarrow$ We have for every $\omega \in A^*$ that the function $\bar{\omega} \circ f$ is analytic $\Leftrightarrow$ We have for every $a \in A$ that the function $O \rightarrow A : z \mapsto f(z)a$ is analytic.*

It is possible to give the following definition.

Definition 1.23 *Consider a complex number z . Then the mapping α_z is closable for the strict topology on $M(A)$ and we denote the strict closure by $\bar{\alpha}_z$.*

In the case where z belongs to $\mathbb{R}$, the previous definition implies that $\bar{\alpha}_z$ is the unique $*$ -homomorphism from $M(A)$ into $M(A)$ which extends α_z . It is also clear that $\alpha_z \subseteq \bar{\alpha}_z$ for every $z \in \mathbb{C}$.

The two previous remarks justify the following notation. For every $z \in \mathbb{C}$ and $a \in D(\bar{\alpha}_z)$, we put $\alpha_z(a) = \bar{\alpha}_z(a)$.

Result 1.24 *Let a be an element in $M(A)$. Then the function $\mathbb{R} \rightarrow M(A) : t \mapsto \alpha_t(a)$ is strictly continuous.*

The following theorem is one of the most important results of [9].

Theorem 1.25 *Consider $z \in \mathbb{C}$. Then we have the following properties*

- *Let a be an element in $M(A)$.*

Then a belongs to $D(\bar{\alpha}_z) \Leftrightarrow$ there exists a function f from $S(z)$ into $M(A)$ such that

1. *f is strictly continuous on $S(z)$*
2. *f is analytic on $S(z)^0$*
3. *We have that $\alpha_t(a) = f(t)$ for every $t \in \mathbb{R}$*

- Let a be an element in $D(\overline{\alpha}_z)$ and f the function from $S(z)$ into $M(A)$ such that

1. f is strictly continuous on $S(z)$
2. f is analytic on $S(z)^0$
3. We have that $\alpha_t(a) = f(t)$ for every $t \in \mathbb{R}$

Then we have that $\alpha_z(a) = f(z)$.

It is also possible to give another characterization of $\overline{\alpha}_z$. First look at the following result.

Proposition 1.26 Consider $z \in \mathbb{C}$ and $a \in D(\overline{\alpha}_z)$, $b \in D(\alpha_z)$. Then ab and ba belong to $D(\alpha_z)$ and $\alpha_z(ab) = \alpha_z(a)\alpha_z(b)$ and $\alpha_z(ba) = \alpha_z(b)\alpha_z(a)$.

There is also a converse of this :

Proposition 1.27 Consider elements $a, b \in M(A)$ and $z \in \mathbb{C}$. Then

1. If we have for every $c \in D(\alpha_z)$ that ac belongs to $D(\alpha_z)$ and $\alpha_z(ac) = b\alpha_z(c)$, then a belongs to $D(\overline{\alpha}_z)$ and $\alpha_z(a) = b$.
2. If we have for every $c \in D(\alpha_z)$ that ca belongs to $D(\alpha_z)$ and $\alpha_z(ca) = \alpha_z(c)b$, then a belongs to $D(\overline{\alpha}_z)$ and $\alpha_z(a) = b$.

These strictly analytic extensions satisfy the same kind of properties as the norm analytic extensions. Now we give a list of these properties.

Remark 1.28 • Consider $z \in \mathbb{C}$ and $y \in S(z)$. Then it is clear that $D(\overline{\alpha}_z)$ is a subset of $D(\overline{\alpha}_y)$.

- It is also clear that $D(\overline{\alpha}_y) = D(\overline{\alpha}_z)$ for $y, z \in \mathbb{C}$ with $\text{Im } y = \text{Im } z$.
- Let $z \in \mathbb{C}$ and $a \in D(\overline{\alpha}_z)$. Then the function $S(z) \rightarrow M(A) : u \mapsto \alpha_u(a)$ is strictly continuous on $S(z)$ and analytic on $S(z)^0$.
- Consider an element a in $M(A)$. We say that a is strictly analytic with respect to α if a belongs to $D(\overline{\alpha}_z)$ for every $z \in \mathbb{C}$. If a is strictly analytic with respect to α , then the function $\mathbb{C} \rightarrow M(A) : u \mapsto \alpha_u(a)$ is analytic.

Proposition 1.29 The mapping $\overline{\alpha}_z$ is a strictly closed multiplicative linear operator in $M(A)$.

Proposition 1.30 Consider $z \in \mathbb{C}$. Then $D(\overline{\alpha}_{\overline{z}}) = D(\overline{\alpha}_z)^*$ and $\text{Ran } \overline{\alpha}_{\overline{z}} = (\text{Ran } \overline{\alpha}_z)^*$. We have for every $a \in D(\alpha_z)$ that $\alpha_z(a)^* = \alpha_{\overline{z}}(a^*)$.

This implies that for all $z \in \mathbb{R}i$ and $a \in D(\overline{\alpha}_z)$ that $\alpha_z(a)^*$ belongs to $D(\overline{\alpha}_z)$ and $\alpha_z(\alpha_z(a)^*)^* = a$.

Result 1.31 Consider $z \in \mathbb{C}$ and $a \in D(\overline{\alpha}_z)$. Then the function $S(z) \rightarrow M(A) : u \mapsto \alpha_u(a)$ is bounded.

Proposition 1.32 Consider $z \in \mathbb{C}$ and $t \in \mathbb{R}$. Then $\overline{\alpha}_z \overline{\alpha}_t = \overline{\alpha}_t \overline{\alpha}_z = \overline{\alpha}_{z+t}$. Furthermore the following equalities hold :

- We have that $\overline{\alpha}_t(D(\overline{\alpha}_z)) = D(\overline{\alpha}_z)$ and $\overline{\alpha}_t(\text{Ran } \overline{\alpha}_z) = \text{Ran } \overline{\alpha}_z$.
- We have that $D(\overline{\alpha}_{z+t}) = D(\overline{\alpha}_z)$ and $\text{Ran } \overline{\alpha}_{z+t} = \text{Ran } \overline{\alpha}_z$.

Proposition 1.33 *Consider $z \in \mathbb{C}$. Then $\overline{\alpha}_z$ is injective and $(\overline{\alpha}_z)^{-1} = \overline{\alpha}_{-z}$. Therefore $\text{Ran } \overline{\alpha}_z = D(\overline{\alpha}_{-z})$ and $D(\overline{\alpha}_z) = \text{Ran } \overline{\alpha}_{-z}$.*

Proposition 1.34 *Consider $y, z \in \mathbb{C}$. Then*

1. $\overline{\alpha}_y \overline{\alpha}_z \subseteq \overline{\alpha}_{y+z}$.
2. *If y and z lie at the same side of the real axis, we have that $\overline{\alpha}_y \overline{\alpha}_z = \overline{\alpha}_{y+z}$.*

As before, we smear elements to construct elements which behave well with respect to α .

Notation 1.35 *Consider $a \in M(A)$, $n > 0$ and $z \in \mathbb{C}$. Then we define the element $a(n, z)$ in $M(A)$ such that*

$$a(n, z)b = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \alpha_t(a) b \, dt$$

for every $b \in A$. Then we have that $\|a(n, z)\| \leq \|a\| \exp(n^2(\text{Im}z)^2)$.

For $n > 0$ and $z \in \mathbb{C}$, we will use the notation $a(n) = a(n, 0)$, so

$$a(n)b = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) b \, dt$$

for every $b \in A$.

Proposition 1.36 *Let $a \in M(A)$, $n > 0$ and $z \in \mathbb{C}$. Then $a(n, z)$ is strictly analytic with respect to α and $\alpha_y(a(n, z)) = a(n, z + y)$ for every $y \in \mathbb{C}$.*

Proposition 1.37 *Consider $n > 0$, $y, z \in \mathbb{C}$ and $a \in D(\overline{\alpha}_y)$. Then $\alpha_y(a(n, z)) = \alpha_y(a)(n, z)$.*

Another useful property of smearing is contained in the following result.

Proposition 1.38 *Let $z \in \mathbb{C}$ and $n > 0$. Consider $a \in M(A)$ and a bounded net $(a_i)_{i \in I}$ in $M(A)$ such that $(a_i)_{i \in I}$ converges strictly to a . Then $(a_i(n, z))_{i \in I}$ is also bounded and converges strictly to $a(n, z)$.*

2 The definition of KMS-weights

In this section, we introduce the definition of a KMS-weight. We will also introduce the definition of a so-called regular weight. In a later section, it is shown that a KMS-weight is automatically regular. We will also prove a result about such a regular weight which will be very useful in a later section. Also some results about lower semi-continuous weights will be included. The standard reference for lower semi-continuous weights is [3].

Let us first introduce notations and conventions concerning weights.

Consider a C^* -algebra A and a densely defined weight φ on A . We use the following notations :

- $\mathcal{M}_\varphi^+ = \{ a \in A^+ \mid \varphi(a) < \infty \}$
- $\mathcal{N}_\varphi = \{ a \in A \mid \varphi(a^*a) < \infty \}$
- $\mathcal{M}_\varphi = \text{span } \mathcal{M}_\varphi^+ = \mathcal{N}_\varphi^* \mathcal{N}_\varphi$.

A GNS-construction of φ is by definition a triple (H, π, Λ) such that

- H is a Hilbert space
- Λ is a linear map from $\mathcal{N}_\varphi$ into H such that
 1. $\Lambda(\mathcal{N}_\varphi)$ is dense in H
 2. We have that $\langle \Lambda(a), \Lambda(b) \rangle = \varphi(b^*a)$ for every $a, b \in \mathcal{N}_\varphi$.
- π is a representation of A on H such that $\pi(a) \Lambda(b) = \Lambda(ab)$ for every $a \in A$ and $b \in \mathcal{N}_\varphi$.

The following concepts play a central role in the theory of lower semi-continuous weights :

- We define the set $\mathcal{F}_\varphi = \{ \omega \in A_+^* \mid \omega \leq \varphi \}$.
- Put $\mathcal{G}_\varphi = \{ \alpha \omega \mid \omega \in \mathcal{F}_\varphi, \alpha \in]0, 1[\}$. Then $\mathcal{G}_\varphi$ is a directed subset of $\mathcal{F}_\varphi$.

A proof of the last result can be found in proposition 3.5.

The major result about lower semi-continuous weights was proven by Combes (proposition 1.7 of [3]) :

Theorem 2.1 *Suppose that φ is lower semi-continuous, then we have the following results :*

- *We have for every $x \in A^+$ that $\varphi(x) = \sup \{ \omega(x) \mid \omega \in \mathcal{F}_\varphi \}$*
- *Consider $x \in A^+$, then the net $(\omega(x))_{\omega \in \mathcal{G}_\varphi}$ converges to $\varphi(x)$*
- *Consider $x \in \mathcal{M}_\varphi$, then the net $(\omega(x))_{\omega \in \mathcal{G}_\varphi}$ converges to $\varphi(x)$*

By proposition 2.4 of [3], we have the following result.

Result 2.2 *Consider $\omega \in \mathcal{F}_\varphi$. Then there exists a unique element $T_\omega \in B(H)$ with $0 \leq T_\omega \leq 1$ such that $\langle T_\omega \Lambda(a), \Lambda(b) \rangle$ for every $a, b \in \mathcal{N}_\varphi$.*

The theorem above implies that $(T_\omega)_{\omega \in \mathcal{G}_\varphi}$ converges strongly to 1 if φ is lower semi-continuous.

Result 2.3 *Suppose that φ is lower semi-continuous. Then we have the following properties.*

1. The mapping Λ is closed.
2. The $*$ -representation π is non-degenerate and $\pi(a)\Lambda(b) = \Lambda(ab)$ for every $a \in M(A)$ and $b \in \mathcal{N}_\varphi$.

Proof :

1. Choose a sequence $(a_n)_{n=1}^\infty \in \mathcal{N}_\varphi$, $a \in A$ and $v \in H$ such that $(a_n)_{n=1}^\infty$ converges to a and $(\Lambda(a_n))_{n=1}^\infty$ converges to v .

Choose $\omega \in \mathcal{F}_\varphi$. We have for every $n \in \mathbb{N}$ that $\langle T_\omega \Lambda(a_n), \Lambda(a_n) \rangle = \omega(a_n^* a_n)$, which implies that $\langle T_\omega v, v \rangle = \omega(a^* a)$. So we see that $\omega(a^* a) \leq \|v\|^2$.

Hence, theorem 2.1 implies that a belongs to $\mathcal{N}_\varphi$.

Choose $\omega \in \mathcal{F}_\varphi$.

Take $b \in \mathcal{N}_\varphi$. We have for every $n \in \mathbb{N}$ that $\langle T_\omega \Lambda(a_n), \Lambda(b) \rangle = \omega(b^* a_n)$, which implies that

$$\langle T_\omega v, \Lambda(b) \rangle = \omega(b^* a) = \langle T_\omega \Lambda(a), \Lambda(b) \rangle .$$

This implies that $T_\omega v = T_\omega \Lambda(a)$.

The lower semi-continuity of φ implies that $(T_\omega)_{\omega \in \mathcal{G}_\varphi}$ converges to 1. Hence $v = \Lambda(a)$.

2. From [3], we know already that π is non-degenerate. Choose $a \in M(A)$ and $b \in \mathcal{N}_\varphi$.

Take $e \in A$. Then ea belongs to A which implies that $ea b$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(eab) = \pi(ea)\Lambda(b) = \pi(e)\pi(a)\Lambda(b)$.

It is clear that $\mathcal{N}_\varphi$ is a left ideal in $M(A)$ so ab belongs to $\mathcal{N}_\varphi$. This implies that $ea b$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(eab) = \pi(e)\Lambda(ab)$.

So we see that $\pi(e)\pi(a)\Lambda(b) = \pi(e)\Lambda(ab)$.

Therefore, the non-degeneracy of π implies that $\Lambda(ab) = \pi(a)\Lambda(b)$.

■

If φ is lower semi-continuous, the weight φ has a natural extension to a weight $\overline{\varphi}$ on $M(A)$ by putting

$$\overline{\varphi}(x) = \sup \{ \omega(x) \mid \omega \in \mathcal{F}_\varphi \}$$

for every $x \in M(A)^+$.

Then $\overline{\varphi}$ is the unique strictly lower semi-continuous weight on $M(A)$ which extends φ . The unicity follows from the fact that any element in $M(A)^+$ can be strictly approximated by an increasing net in A^+ .

We define $\overline{\mathcal{M}}_\varphi = \mathcal{M}_{\overline{\varphi}}$ and $\overline{\mathcal{N}}_\varphi = \mathcal{N}_{\overline{\varphi}}$. For any $x \in \overline{\mathcal{M}}_\varphi$, we put $\varphi(x) = \overline{\varphi}(x)$.

Theorem 2.1 implies that $\overline{\varphi}$ is an extension of φ . So we have that $\mathcal{M}_\varphi^+ = \overline{\mathcal{M}}_\varphi^+ \cap A$, $\mathcal{N}_\varphi = \overline{\mathcal{N}}_\varphi \cap A$ and $\mathcal{M}_\varphi \subseteq \overline{\mathcal{M}}_\varphi \cap A$

Then we have that

- Consider $x \in M(A)^+$, then the net $(\omega(x))_{\omega \in \mathcal{G}_\varphi}$ converges to $\varphi(x)$
- Consider $x \in \overline{\mathcal{M}}_\varphi$, then the net $(\omega(x))_{\omega \in \mathcal{G}_\varphi}$ converges to $\varphi(x)$

Lemma 2.4 Suppose that φ is lower semi-continuous. Then the mapping $\Lambda : \mathcal{N}_\varphi \mapsto H$ is strictly closable.

Proof : Choose a net $(a_i)_{i \in I}$ in $\mathcal{N}_\varphi$ and $v \in H$ such that $(a_i)_{i \in I}$ converges strictly to 0 and such that $(\Lambda(a_i))_{i \in I}$ converges to v .

Take $e \in A$. Then we have that $(ea_i)_{i \in I}$ converges to 0. We have also for every $i \in I$ that ea_i belongs to $\mathcal{N}_\varphi$ and $\Lambda(ea_i) = \pi(e)\Lambda(a_i)$. Hence, $(\Lambda(ea_i))_{i \in I}$ converges to $\pi(e)v$. So the closedness of Λ implies that $\pi(e)v = 0$.

Therefore, the non-degeneracy of π implies that $v = 0$. ■

Definition 2.5 Suppose that φ is lower semi-continuous. Then we define the mapping $\overline{\Lambda}$ from within $M(A)$ into H such that $\mathcal{N}_\varphi$ is a strict core for $\overline{\Lambda}$ and $\overline{\Lambda}$ extends Λ . We put $\Lambda(a) = \overline{\Lambda}(a)$ for every $a \in D(\overline{\Lambda})$

Using result 2.3.2, it is not very difficult to prove for every $x \in M(A)$ and every $a \in D(\overline{\Lambda})$ that xa belongs to $D(\overline{\Lambda})$ and $\Lambda(xa) = \pi(a)\Lambda(x)$.

Proposition 2.6 Suppose that φ is lower semi-continuous. Then $(H, \overline{\Lambda}, \overline{\pi})$ is a GNS-construction for $\overline{\varphi}$.

Proof :

- Choose $a \in D(\overline{\Lambda})$. By definition, there exists a net $(a_i)_{i \in I}$ in $\mathcal{N}_\varphi$ such that $(a_i)_{i \in I}$ converges strictly to a and $(\Lambda(a_i))_{i \in I}$ converges to $\Lambda(a)$.

Take $\omega \in \mathcal{F}_\varphi$. Then we have for every $i, j \in I$ that $\langle T_\omega \Lambda(a_i), \Lambda(a_j) \rangle = \omega(a_j^* a_i)$.

This implies that $\langle T_\omega \Lambda(a), \Lambda(a) \rangle = \omega(a^* a)$.

By the definition of $\overline{\varphi}$ and because $(T_\omega)_{\omega \in \mathcal{G}_\varphi}$ converges strongly to 1, we get that a belongs to $\overline{\mathcal{N}}_\varphi$ and that $\varphi(a^* a) = \langle \Lambda(a), \Lambda(a) \rangle$.

- Choose $a \in \overline{\mathcal{N}}_\varphi$. Take an approximate unit $(e_i)_{i \in I}$ of A . Then $(e_i a)_{i \in I}$ converges strictly to a .

We have for every $i \in I$ that $e_i a \in \overline{\mathcal{N}}_\varphi \cap A = \mathcal{N}_\varphi$.

Because $(a^* e_i a)_{i \in I}$ is an increasing net in $\overline{\mathcal{M}}_\varphi^+$, which converges strictly to $a^* a$, the strict lower semi-continuity of $\overline{\varphi}$ implies that $(\varphi(a^* e_i a))_{i \in I}$ converges to $\varphi(a^* a)$.

We have moreover for every $j, k \in I$ with $j \leq k$ that $0 \leq e_k - e_j \leq 1$, so

$$\|\Lambda(e_j a) - \Lambda(e_k a)\| = \varphi(a^* (e_k - e_j)^2 a) \leq \varphi(a^* (e_k - e_j) a) = \varphi(a^* e_k a) - \varphi(a^* e_j a)$$

So we get that $(\Lambda(e_i a))_{i \in I}$ is cauchy and hence convergent in H .

Hence, we get by the definition of $\overline{\Lambda}$ that a belongs to $D(\overline{\Lambda})$. ■

Remark 2.7 By definition, we have that $\mathcal{N}_\varphi$ is a strict core for $\overline{\Lambda}$. But we have even more: Consider $a \in \overline{\mathcal{N}}_\varphi$. Then there exists a net $(a_k)_{k \in K}$ in $\mathcal{N}_\varphi$ such that

- We have for every $k \in K$ that $\|a_k\| \leq \|a\|$ and $\|\Lambda(a_k)\| \leq \|\Lambda(a)\|$.

- The net $(a_k)_{k \in K}$ converges strictly to a and the net $(\Lambda(a_k))_{k \in K}$ converges strongly* to $\Lambda(a)$.

This follows immediately by multiplying a to the left by an approximate unit of A .

We will not use this extension to the multiplier in this paper. However, most of the results in this paper concerning φ , $\mathcal{M}_\varphi$, $\mathcal{M}_\varphi^+$, $\mathcal{N}_\varphi$, Λ have obvious variants concerning φ , $\overline{\mathcal{M}}_\varphi^+$, $\overline{\mathcal{M}}_\varphi$, $\overline{\mathcal{N}}_\varphi$, $\overline{\Lambda}$. They can be proven using the results in this paper (using strict approximation arguments) or by similar proofs.

Now we give the definition of a KMS-weight.

Definition 2.8 *Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A such that there exist a norm-continuous one parameter group σ on A such that*

1. *We have that $\varphi \sigma_t = \varphi$ for every $t \in \mathbb{R}$.*
2. *We have that $\varphi(a^*a) = \varphi(\sigma_{\frac{i}{2}}(a) \sigma_{\frac{i}{2}}(a)^*)$ for every $a \in D(\sigma_{\frac{i}{2}})$.*

Then φ is called a KMS-weight and σ is called a modular group for φ .

Later we will prove that the second condition on σ can be weakened.

This definition of a KMS-weight is different from the usual one (see definition 4.1 of [4]) but we will prove in theorem 6.36 that this definition is equivalent with the usual one.

In [16], Jan Verding introduced the notion of so-called regular weights. We will formulate this definition and prove a useful property concerning regular weights.

Definition 2.9 *Consider a C^* -algebra A . Let φ be a weight on A with GNS-construction (H, Λ, π) . We call φ regular $\Leftrightarrow \varphi$ is densely defined, lower semi-continuous and there exists a net $(u_i)_{i \in I}$ in $\mathcal{N}_\varphi$ such that*

- *We have for every $i \in I$ that :*
 1. *There exists $S_i \in B(H)$ such that $S_i \Lambda(a) = \Lambda(a u_i)$ for every $a \in \mathcal{N}_\varphi$.*
 2. *$\|u_i\| \leq 1$ and $\|S_i\| \leq 1$.*
- *Furthermore, we assume that*
 1. *$(u_i)_{i \in I}$ converges strictly to 1.*
 2. *$(S_i)_{i \in I}$ converges strongly to 1.*

We call the net $(u_i)_{i \in I}$ a truncating net for φ .

It is not difficult to see that the definition of regularity is independent of the choice of GNS-construction.

Proposition 2.10 *Consider a regular weight φ on a C^* -algebra A with GNS-construction (H, Λ, π) . Let B be a sub- $*$ -algebra of $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ such that B is a core for Λ and let a be an element in $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. Then there exists a sequence $(a_n)_{n=1}^\infty$ in B such that*

1. $(a_n)_{n=1}^\infty \rightarrow a$
2. $(\Lambda(a_n))_{n=1}^\infty \rightarrow \Lambda(a)$

$$3. (\Lambda(a_n^*))_{n=1}^\infty \rightarrow \Lambda(a^*).$$

Proof : Choose a truncating net $(u_i)_{i \in I}$ for φ . Define for every $i \in I$ the operator $S_i \in B(H)$ such that $S_i \Lambda(a) = \Lambda(au_i)$ for every $a \in \mathcal{N}_\varphi$.

Choose $n \in \mathbb{N}$. Because $(u_i)_{i \in I}$ is a bounded net which converges strictly to 1 and $(S_i)_{i \in I}$ is a bounded net which converges strongly to 1, there exist an element $j \in I$ such that

- $\|\Lambda(a) - \pi(u_j^*)S_j\Lambda(a)\| \leq \frac{1}{n}$
- $\|\Lambda(a^*) - \pi(u_j^*)S_j\Lambda(a^*)\| \leq \frac{1}{n}$
- $\|a - u_j^*au_j\| \leq \frac{1}{n}$.

By the definition of S_j , we get that

- $\|\Lambda(a) - \pi(u_j^*)\pi(a)\Lambda(u_j)\| \leq \frac{1}{n}$
- $\|\Lambda(a^*) - \pi(u_j^*)\pi(a^*)\Lambda(u_j)\| \leq \frac{1}{n}$.

Choose $n \in \mathbb{N}$. Because B is a core for Λ , there exists an element $b_n \in B$ such that

- $\|u_j - b_n\| \leq \min\{\frac{1}{2n} \frac{1}{\|a\|+1}, \frac{1}{n} \frac{1}{(\|a\|+1)(\|\Lambda(u_j)\|+1)}\}$
- $\|\Lambda(u_j) - \Lambda(b_n)\| \leq \frac{1}{2n} \frac{1}{\|a\|+1}$
- $\|b_n\| \leq 2$

Furthermore, there exists an element $d_n \in B$ such that

$$\|a - d_n\| \leq \min\{\frac{1}{n} \frac{1}{\|b_n\|^2 + 1}, \frac{1}{n} \frac{1}{(\|b_n\| + 1)(\|\Lambda(b_n)\| + 1)}\}$$

We define $a_n = b_n^*d_nb_n$. Then a_n belongs to B and $a_n^* = b_n^*d_n^*b_n$.

Moreover,

1. Using the above estimations, we have that

$$\begin{aligned} \|\Lambda(a_n) - \Lambda(a)\| &= \|\pi(b_n^*)\pi(d_n)\Lambda(b_n) - \Lambda(a)\| \\ &\leq \|\pi(b_n^*)\pi(d_n)\Lambda(b_n) - \pi(b_n^*)\pi(a)\Lambda(b_n)\| + \|\pi(b_n^*)\pi(a)\Lambda(b_n) - \pi(b_n^*)\pi(a)\Lambda(u_j)\| \\ &\quad + \|\pi(b_n^*)\pi(a)\Lambda(u_j) - \pi(u_j^*)\pi(a)\Lambda(u_j)\| + \|\pi(u_j^*)\pi(a)\Lambda(u_j) - \Lambda(a)\| \\ &\leq \|b_n\| \|\Lambda(b_n)\| \|d_n - a\| + \|b_n\| \|a\| \|\Lambda(b_n) - \Lambda(u_j)\| + \|b_n - u_j\| \|a\| \|\Lambda(u_j)\| + \frac{1}{n} \\ &\leq \frac{1}{n} + 2 \|a\| \|\Lambda(b_n) - \Lambda(u_j)\| + \frac{1}{n} + \frac{1}{n} \leq \frac{1}{n} + \frac{1}{n} + \frac{1}{n} + \frac{1}{n} = \frac{4}{n} \end{aligned}$$

2. Similarly, one can prove $\|\Lambda(a_n^*) - \Lambda(a^*)\| \leq \frac{4}{n}$.

3. Using the above estimations once again, we get that

$$\begin{aligned} \|a_n - a\| &= \|b_n^*d_nb_n - a\| \\ &\leq \|b_n^*d_nb_n - b_n^*ab_n\| + \|b_n^*ab_n - b_n^*au_j\| + \|b_n^*au_j - u_j^*au_j\| + \|u_j^*au_j - a\| \\ &\leq \|b_n\|^2 \|d_n - a\| + \|b_n\| \|a\| \|b_n - u_j\| + \|b_n - u_j\| \|a\| \|u_j\| + \|u_j^*au_j - a\| \\ &\leq \frac{1}{n} + 2 \|a\| \|b_n - u_j\| + \|b_n - u_j\| \|a\| + \frac{1}{n} \leq \frac{1}{n} + \frac{1}{n} + \frac{1}{n} + \frac{1}{n} = \frac{4}{n}. \end{aligned}$$

Looking at 1, 2 and 3, we see that $(a_n)_{n=1}^\infty \rightarrow a$, $(\Lambda(a_n))_{n=1}^\infty \rightarrow \Lambda(a)$ and $(\Lambda(a_n^*))_{n=1}^\infty \rightarrow \Lambda(a^*)$. ■

Corollary 2.11 *Consider a regular weight φ on a C^* -algebra A with GNS-construction (H, Λ, π) . Let a be an element in $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. Then there exists a sequence $(a_n)_{n=1}^\infty$ in $\mathcal{M}_\varphi$ such that*

1. $(a_n)_{n=1}^\infty \rightarrow a$
2. $(\Lambda(a_n))_{n=1}^\infty \rightarrow \Lambda(a)$
3. $(\Lambda(a_n^*))_{n=1}^\infty \rightarrow \Lambda(a^*)$.

We need the following result from [14] (lemma 3.1). The proof remains valid for lower semi-continuous weights on C^* -algebras.

Lemma 2.12 *Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A . Let f be a function from $\mathbb{R}$ into $\mathcal{M}_\varphi^+$ which is continuous and integrable. Then the function $\mathbb{R} \rightarrow \mathbb{R}^+ : t \mapsto \varphi(f(t))$ is lower semi-continuous and*

$$\varphi\left(\int f(t) dt\right) = \int \varphi(f(t)) dt .$$

Lemma 2.13 *Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A , α a norm continuous one-parameter group on A such that there exists a strictly positive number λ such that $\varphi \alpha_t = \lambda^t \varphi$ for every $t \in \mathbb{R}$. Let $z \in \mathbb{C}$, $n \in \mathbb{N}$ and $a \in \mathcal{M}_\varphi$.*

Then we have that the element $\int \exp(-(t-z)^2) \alpha_t(a) dt$ belongs to $\mathcal{M}_\varphi$ and

$$\varphi\left(\int \exp(-(t-z)^2) \alpha_t(a) dt\right) = \lambda^z \sqrt{\pi} \exp\left(\frac{(\ln \lambda)^2}{4}\right) \varphi(a) .$$

Proof : Choose $b \in \mathcal{M}_\varphi^+$. Define the function $f : \mathbb{R} \rightarrow \mathbb{C} : t \mapsto \exp(-(t-z)^2)$.

There exist continuous positive functions $f_1, \dots, f_4$ such that $f_1, \dots, f_4 \leq |f|$ and $f = f_1 - f_2 + if_3 - if_4$.

Fix $j \in \{1, \dots, 4\}$. It is clear that the function $\mathbb{R} \rightarrow \mathcal{M}_\varphi^+ : t \mapsto f_j(t) \alpha_t(b)$ is continuous. Because we also have that $\|f_j(t) \alpha_t(b)\| = f_j(t) \|b\| \leq |f(t)| \|b\|$ for every $t \in \mathbb{R}$, we see that the function $\mathbb{R} \rightarrow \mathcal{M}_\varphi^+ : t \mapsto f_j(t) \alpha_t(b)$ is integrable. So, the previous lemma implies that the function $\mathbb{R} \rightarrow \mathbb{R}^+ : t \mapsto \varphi(f_j(t) \alpha_t(b))$ is lower semi-continuous and

$$\varphi\left(\int f_j(t) \alpha_t(b) dt\right) = \int \varphi(f_j(t) \alpha_t(b)) dt = \left(\int f_j(t) \lambda^t dt\right) \varphi(b) .$$

Because $f_j \leq |f|$, the function $\mathbb{R} \rightarrow \mathbb{R}^+ : t \mapsto f_j(t) \lambda^t$ is integrable. Therefore we see that $\int f_j(t) \alpha_t(b) dt$ belongs to $\mathcal{M}_\varphi^+$.

Using the fact that $f = f_1 - f_2 + if_3 - if_4$, we get that $\int f(t) \alpha_t(b) dt$ belongs to $\mathcal{M}_\varphi$ and

$$\varphi\left(\int f(t) \alpha_t(b) dt\right) = \left(\int f(t) \lambda^t dt\right) \varphi(b) .$$

Therefore,

$$\begin{aligned} \varphi\left(\int f(t) \alpha_t(b) dt\right) &= \left(\int \exp(-(t-z)^2) \lambda^t dt\right) \varphi(b) \\ &= \left(\int \exp(-t^2) \lambda^{t+z} dt\right) \varphi(b) = \lambda^z \sqrt{\pi} \exp\left(\frac{(\ln \lambda)^2}{4}\right) \varphi(b) . \end{aligned}$$

The lemma follows by linearity. ■

Proposition 2.14 Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A , α a norm continuous one-parameter group on A such that there exists a strictly positive number λ such that $\varphi \alpha_t = \lambda^t \varphi$ for every $t \in \mathbb{R}$. Let $z \in \mathbb{C}$ and $a \in \mathcal{M}_\varphi \cap D(\alpha_z)$ such that $\alpha_z(a)$ belongs to $\mathcal{M}_\varphi$. Then we have that $\varphi(\alpha_z(a)) = \lambda^z \varphi(a)$.

Proof : Define

$$x = \int \exp(-t^2) \alpha_t(\alpha_z(a)) dt .$$

Because $\alpha_z(a)$ belongs to $\mathcal{M}_\varphi$, the previous lemma implies that x belongs to $\mathcal{M}_\varphi$ and

$$\varphi(x) = \sqrt{\pi} \exp\left(\frac{(\ln \lambda)^2}{4}\right) \varphi(\alpha_z(a)) .$$

We have also that

$$x = \int \exp(-(t-z)^2) \alpha_t(a) dt ,$$

so the previous lemma implies in this case that

$$\varphi(x) = \lambda^z \sqrt{\pi} \exp\left(\frac{(\ln \lambda)^2}{4}\right) \varphi(a) .$$

Comparing these two different expressions for $\varphi(x)$ gives us that $\varphi(\alpha_z(a)) = \lambda^z \varphi(a)$. ■

Result 2.15 Consider a densely defined lower semi-continuous non-zero weight φ on a C^* -algebra A . Let α be a norm continuous one parameter group on A such that there exists for every t in $\mathbb{R}$ a number λ_t such that $\varphi \alpha_t = \lambda_t \varphi$ for every $t \in \mathbb{R}$. Then there exists a unique strictly positive number λ such that $\varphi \alpha_t = \lambda^t \varphi$ for every $t \in \mathbb{R}$.

Proof : The technique of this proof comes from proposition 5.7 of [14] but can be more easily applied in this case. We have certainly that $\lambda_t > 0$ for every $t \in \mathbb{R}$. Put $\lambda = \lambda_1 > 0$.

Because $\alpha_{s+t} = \alpha_s \alpha_t$ for every $s, t \in \mathbb{R}$, we get easily that $\lambda_{s+t} = \lambda_s \lambda_t$ for every $s, t \in \mathbb{R}$. As usual, this implies that $\lambda_q = \lambda^q$ for every $q \in \mathbb{Q}$.

The lower semi-continuity of φ implies that the mapping $\mathbb{R} \rightarrow \mathbb{R}_0^+ : t \mapsto \lambda_t$ is lower semi continuous. So the mapping $\mathbb{R} \rightarrow \mathbb{R} : t \mapsto \lambda_t - \lambda^t$ is lower semi-continuous. This implies that the set $\{t \in \mathbb{R} \mid \lambda_t \leq \lambda^t\}$ is closed in $\mathbb{R}$. We know already that this set contains $\mathbb{Q}$, so it must be equal to $\mathbb{R}$.

Therefore $\lambda_t \leq \lambda^t$ for every $t \in \mathbb{R}$. This implies that $\lambda_t = \lambda^t$ for every $t \in \mathbb{R}$ (if there would exist an element in $\mathbb{R}$ for which this equality does not hold, this element or its opposite would violate the previous inequality). ■

The proof of the following result is due to J. Verding (see [22]).

Result 2.16 Consider a C^* -algebra A and a densely defined lower semi-continuous weight φ on A . Take a GNS-construction (H, Λ, π) for φ . Let α be a norm continuous one parameter group on A such that there exists a strictly positive number λ such that $\varphi \alpha_t = \lambda^t \varphi$ for every $t \in \mathbb{R}$. Then there exists a unique injective positive operator T in H such that $T^{it} \Lambda(a) = \lambda^{-\frac{t}{2}} \Lambda(\alpha_t(a))$ for every $a \in \mathcal{N}_\varphi$ and $t \in \mathbb{R}$.

Proof : For every $t \in \mathbb{R}$, there exist a unitary operator u_t on H such that $u_t \Lambda(a) = \lambda^{-\frac{t}{2}} \Lambda(\alpha_t(a))$ for every $a \in \mathcal{N}_\varphi$. The mapping $\mathbb{R} \rightarrow B(H) : t \mapsto u_t$ is clearly a unitary representation of $\mathbb{R}$ on H

Fix $\omega \in \mathcal{G}_\varphi$ for the moment. Because $\omega \leq \varphi$, there exists a unique element $T_\omega \in B(H)^+$ such that $\langle T_\omega \Lambda(a), \Lambda(b) \rangle = \omega(b^*a)$ for every $a, b \in \mathcal{N}_\varphi$ (see result 2.2).

Because φ is lower semi-continuous, the remarks after result 2.2 imply that $(T_\omega)_{\omega \in \mathcal{G}_\varphi}$ converges strongly to 1. This implies that the set $\{T_\omega \Lambda(a) \mid \omega \in \mathcal{G}_\varphi, a \in \mathcal{N}_\varphi\}$ is dense in H . (*)

Choose $\theta \in \mathcal{G}_\varphi$ and $x, y \in \mathcal{N}_\varphi$. We have for every $t \in \mathbb{R}$ that

$$\langle u_t \Lambda(x), T_\theta \Lambda(y) \rangle = \lambda^{-\frac{t}{2}} \langle \Lambda(\alpha_t(x)), T_\theta \Lambda(y) \rangle = \lambda^{-\frac{t}{2}} \theta(y^* \alpha_t(x)) .$$

This implies that the mapping $\mathbb{R} \rightarrow \mathbb{C} : t \mapsto \langle u_t \Lambda(x), T_\theta \Lambda(y) \rangle$ is continuous.

Therefore (and because $\|u_t\| \leq 1$ for every $t \in \mathbb{R}$), result (*) implies that the mapping $\mathbb{R} \rightarrow \mathbb{C} : t \mapsto \langle u_t v, w \rangle$ is continuous for all $v, w \in H$.

Consequently, the mapping $\mathbb{R} \rightarrow B(H) : t \mapsto u_t$ is a strongly continuous unitary group representation of $\mathbb{R}$ on H . The result follows by the Stone theorem. ■

Terminology 2.17 Consider a C^* -algebra A and an element T affiliated with A . A truncating sequence for T is by definition a sequence $(e_n)_{n=1}^\infty$ in $M(A)$ such that :

1. $(e_n)_{n=1}^\infty$ is bounded and converges strictly to 1.
2. We have for every $n \in \mathbb{N}$ that e_n is a left and right multiplier of $|T|$ and $|T| e_n = e_n |T|$.

It is not difficult to check that the functional calculus for $|T|$ guarantees the existence of such a truncating sequence for T .

Let us fix for the moment an element T affiliated with a C^* -algebra A and a truncating sequence $(e_n)_{n=1}^\infty$ for T . Then we have the following properties.

- Take $m \in \mathbb{N}$. We know for every $x \in A$ that $e_m x$ belongs to $D(|T|)$, so $e_m x$ belongs to $D(T)$. Furthermore

$$\|T(e_m x)\| = \||T|(e_m x)\| .$$

This implies that $T e_m$ is a bounded linear operator on A with $\|T e_m\| = \||T| e_m\|$. So e_m is a right multiplier of T .

- Let $x \in D(T)$. Then we have for every $n \in \mathbb{N}$ that

$$\|T(e_n x) - T(x)\| = \||T|(e_n x) - |T|(x)\| = \|e_n |T|(x) - |T|(x)\| .$$

This implies that $(T(e_n x))_{n=1}^\infty$ converges to $T(x)$.

Combining this with the closedness of T , this implies for every $x \in A$ that x belongs to $D(T)$ if and only if the sequence $(T(e_n x))_{n=1}^\infty$ is convergent in A .

Proposition 2.18 Consider a C^* -algebra A and an element T affiliated with A . Let φ be a densely defined lower semi-continuous weight on A with GNS-construction (H, Λ, π) . Let a be an element in $D(T) \cap \mathcal{N}_\varphi$, then :

1. We have that $T(a)$ belongs to $\mathcal{N}_\varphi \Leftrightarrow \Lambda(a)$ belongs to $D(\pi(T))$.
2. If $T(a)$ belongs to $\mathcal{N}_\varphi$, then $\Lambda(T(a)) = \pi(T)\Lambda(a)$.

Proof : Take a truncating sequence $(e_n)_{n=1}^\infty$ for T . It is not too difficult to check that $\pi(T)\pi(e_n) = \pi(Te_n) \in B(H)$ and $\pi(e_n)|\pi(T)| \subseteq |\pi(T)|\pi(e_n) = \pi(|T|e_n) \in B(H)$.

- Suppose that $T(a)$ belongs to $\mathcal{N}_\varphi$. Because a belongs to $D(|T|)$ and $(|T|a)^*(|T|a) = (Ta)^*(Ta)$, we see that $|T|a$ also belongs to $\mathcal{N}_\varphi$.

Fix $n \in \mathbb{N}$. We know that $|T|e_n$ belongs to $M(A)$ and $e_n|T| \subseteq |T|e_n$. We know also that $\pi(|T|)\pi(e_n)$ belongs to $B(H)$ and $|\pi(T)|\pi(e_n) = \pi(|T|e_n)$.

Hence, $\pi(e_n)\Lambda(a)$ belongs to $D(|\pi(T)|)$ and

$$\begin{aligned} |\pi(T)|(\pi(e_n)\Lambda(a)) &= \pi(|T|e_n)\Lambda(a) = \Lambda(|T|e_n)a \\ &= \Lambda(e_n(|T|a)) = \pi(e_n)\Lambda(|T|a) . \end{aligned}$$

This last equality implies that $(|\pi(T)|(\pi(e_n)\Lambda(a)))_{n=1}^\infty$ converges to $\Lambda(|T|a)$. Therefore $\Lambda(a)$ must belong to $D(|\pi(T)|)$, so it must belong to $D(\pi(T))$.

- Suppose that $\Lambda(a)$ belongs to $D(\pi(T))$. Then $\Lambda(a)$ will also belong to $D(|\pi(T)|)$.

For every $n \in N$, we have that Te_n belongs to $M(A)$, which implies that $(Te_n)a$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda((Te_n)a) = \pi(Te_n)\Lambda(a) = \pi(T)(\pi(e_n)\Lambda(a)) .$$

Because $\Lambda(a)$ belongs to $D(|\pi(T)|)$, this equality implies for every $n \in \mathbb{N}$ that

$$\begin{aligned} \|\Lambda((Te_n)a) - \pi(T)\Lambda(a)\| &= \|\pi(T)(\pi(e_n)\Lambda(a) - \Lambda(a))\| \\ &= \|\pi(T)|(\pi(e_n)\Lambda(a) - \Lambda(a))\| = \|\pi(e_n)|\pi(T)|\Lambda(a) - |\pi(T)|\Lambda(a)\| \end{aligned}$$

So $(\Lambda((Te_n)a))_{n=1}^\infty$ converges to $\pi(T)\Lambda(a)$.

We also know that $(Te_n a)_{n=1}^\infty$ converges to $T(a)$, so the closedness of Λ implies that $T(a)$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(T(a)) = \pi(T)\Lambda(a)$. ■

3 Weights arising from GNS-constructions

In this section, we gather some results from [22] and [15]. Except lemma 3.3, all the results and proofs in this section are due to Jan Verding and can be found in [22] and [15]. However, due to the lack of availability of these two works, we feel it necessary to include the proofs here.

Consider a C^* -algebra A , a Hilbert space H , a dense left ideal N of A and a linear map Λ from N into H with dense range. Assume furthermore the existence of a $*$ -homomorphism π from A into $B(H)$ such that $\pi(x)\Lambda(a) = \Lambda(xa)$ for every $x \in A$ and $a \in N$.

Notation 3.1 *We define the sets*

$$\mathcal{F} = \{ \omega \in A^+ \mid \text{We have that } \omega(a^*a) \leq \|\Lambda(a)\|^2 \text{ for every } a \in N \}$$

and

$$\mathcal{G} = \{ \lambda \omega \mid \omega \in \mathcal{F}, \lambda \in]0, 1[\} .$$

Remark 3.2 Consider $\omega \in \mathcal{F}$. Then it is not difficult to see that there exists a unique element $T \in B(H)$ with $0 \leq T \leq 1$ such that $\langle T\Lambda(a), \Lambda(b) \rangle = \omega(b^*a)$ for every $a, b \in N$. We get also that T belongs to $\pi(A)'$. The techniques used to prove these results can also be found in the proof of the next lemma.

Lemma 3.3 Consider a positive sesquilinear form on N such that $s(ab_1, b_2) = s(b_1, a^*b_2)$ for all $a \in A$ and all $b_1, b_2 \in N$. Suppose moreover that there exists a positive linear functional θ on A such $s(b, b) \leq \theta(b^*b)$ for every $b \in N$. Then there exists a unique positive linear functional ω on A with $\omega \leq \theta$ such that $\omega(b_2^*b_1) = s(b_1, b_2)$ for every $b_1, b_2 \in N$.

Proof : Let (π, H, v) be GNS-object for θ (v is a cyclic vector).

Because s is a positive sesquilinear form on N , we can use the Cauchy-Schwarz inequality for s . So we have for every $b_1, b_2 \in N$ that

$$|s(b_1, b_2)|^2 \leq s(b_1, b_1) s(b_2, b_2) \leq \theta(b_1^*b_1) \theta(b_2^*b_2) = \|\pi(b_1)v\|^2 \|\pi(b_2)v\|^2.$$

Therefore we can define a continuous positive sesquilinear form t on H such that $t(\pi(b_1)v, \pi(b_2)v) = s(b_1, b_2)$ for every $b_1, b_2 \in N$. It is clear that t is positive and $\|t\| \leq 1$.

So there exist an element $T \in B(H)$ with $0 \leq T \leq 1$ such that $t(x, y) = \langle Tx, y \rangle$ for every $x, y \in H$.

This implies that $\langle T\pi(b_1)v, \pi(b_2)v \rangle = s(b_1, b_2)$ for every $b_1, b_2 \in N$.

Next we show that T belongs to $\pi(A)'$. Therefore, choose $a \in N$.

We have for every $b_1, b_2 \in N$ that

$$\begin{aligned} \langle T\pi(a)\pi(b_1)v, \pi(b_2)v \rangle &= \langle T\pi(ab_1)v, \pi(b_2)v \rangle = s(ab_1, b_2) \\ &= s(b_1, a^*b_2) = \langle T\pi(b_1)v, \pi(a^*b_2)v \rangle \\ &= \langle T\pi(b_1)v, \pi(a^*)\pi(b_2)v \rangle = \langle \pi(a)T\pi(b_1)v, \pi(b_2)v \rangle. \end{aligned}$$

This implies that $T\pi(a) = \pi(a)T$.

Now we define the continuous linear functional ω on A such $\omega(x) = \langle T\pi(x)v, v \rangle$ for every $x \in A$.

Using the fact that T belongs to $\pi(A)'$, we have for every $b_1, b_2 \in N$ that

$$\omega(b_2^*b_1) = \langle T\pi(b_2^*b_1)v, v \rangle = \langle T\pi(b_1)v, \pi(b_2)v \rangle = s(b_1, b_2).$$

Consequently, we have for every $b \in N$ that $\omega(b^*b) = s(b, b)$, implying that $0 \leq \omega(b^*b) \leq \theta(b^*b)$. This implies easily that $0 \leq \omega \leq \theta$. ■

Lemma 3.4 Consider a unital C^* -algebra C . Let T_1, T_2 be elements in C with $0 \leq T_1, T_2 \leq 1$, let γ be a number in $]0, 1[$. Then there exist an element $T \in C$ with $0 \leq T \leq 1$ and such that $\gamma T_1 \leq T$, $\gamma T_2 \leq T$ and $T \leq \frac{\gamma}{1-\gamma}(T_1 + T_2)$.

Proof : For the moment, fix $i \in \{1, 2\}$. We define

$$S_i = \frac{\gamma T_i}{1 - \gamma T_i} \in C.$$

It is then easy to check that

$$\gamma T_i = \frac{S_i}{1 + S_i} \quad (\text{a}) \quad \text{and} \quad S_i \leq \frac{\gamma}{1 - \gamma} T_i \quad (\text{b}).$$

Next, we define

$$T = \frac{S_1 + S_2}{1 + S_1 + S_2} \in C.$$

We get immediately that $0 \leq T \leq 1$. By (b), we have that $T \leq S_1 + S_2 \leq \frac{\gamma}{1-\gamma}(T_1 + T_2)$.

We know that the function $\mathbb{R}^+ \rightarrow \mathbb{R}^+ : t \mapsto \frac{t}{1+t}$ is operator monotone (see [13]). This implies for every $i = 1, 2$ that

$$T \geq \frac{S_i}{S_i + 1} = \gamma T_i,$$

where we used (a) in the last equality. ■

We now use these two lemmas to proof the following useful proposition.

Proposition 3.5 *The set $\mathcal{G}$ is upwardly directed for the natural order on A_+^* .*

Proof : Choose $\omega_1, \omega_2 \in \mathcal{F}$ and $\lambda_1, \lambda_2 \in]0, 1[$. Then there exist a number $\gamma \in]0, 1[$ such that $\lambda_1, \lambda_2 < \gamma$. By the remark after notation 3.1, we know that there exist T_1, T_2 in $\pi(A)'$ such that $0 \leq T_1, T_2 \leq 1$ and such that $\omega_1(y^*x) = \langle T_1 \Lambda(x), \Lambda(y) \rangle$ and $\omega_2(y^*x) = \langle T_2 \Lambda(x), \Lambda(y) \rangle$ for every $x, y \in N$.

The previous lemma implies the existence of $T \in \pi(A)'$ with $0 \leq T \leq 1$ and such that $\gamma T_1 \leq T$, $\gamma T_2 \leq T$ and $T \leq \frac{\gamma}{1-\gamma}(T_1 + T_2)$.

Put $\lambda = \max(\frac{\lambda_1}{\gamma}, \frac{\lambda_2}{\gamma}) \in]0, 1[$. Then $\lambda T \geq \lambda \gamma T_1 \geq \lambda_1 T_1$ and analogously, $\lambda T \geq \lambda_2 T_2$. (a)

Define $\theta = \frac{\gamma}{1-\gamma}(\omega_1 + \omega_2) \in A_+^*$

Next, we define the mapping s from $N \times N$ into $\mathbb{C}$ such that $s(x, y) = \langle T \Lambda(x), \Lambda(y) \rangle$ for every $x, y \in N$. Then s is sesquilinear. Furthermore:

- We have for $x \in N$ that

$$0 \leq \langle T \Lambda(x), \Lambda(y) \rangle \leq \frac{\gamma}{1-\gamma} \langle (T_1 + T_2) \Lambda(x), \Lambda(y) \rangle = \frac{\gamma}{1-\gamma} (\omega_1(x^*x) + \omega_2(x^*x)),$$

which implies that $0 \leq s(x, x) \leq \theta(x^*x)$.

- We have for every $x, y \in N$ and $a \in A$ that

$$\begin{aligned} \langle T \Lambda(ax), \Lambda(y) \rangle &= \langle T \pi(a) \Lambda(x), \Lambda(y) \rangle = \langle \pi(a) T \Lambda(x), \Lambda(y) \rangle \\ &= \langle T \Lambda(x), \pi(a^*) \Lambda(y) \rangle = \langle T \Lambda(x), \Lambda(a^*y) \rangle \end{aligned}$$

which implies that $s(ax, y) = s(x, a^*y)$.

This allows us to apply the previous lemma. Therefore, we get the existence of $\omega \in A_+^*$ with $\omega \leq \theta$ and such that $s(x, y) = \omega(y^*x)$ for every $x, y \in N$.

Then we have for every $x \in N$ that

$$\omega(x^*x) = s(x, x) = \langle T \Lambda(x), \Lambda(x) \rangle \leq \|\Lambda(x)\|^2$$

which implies that ω belongs to $\mathcal{F}$.

By (a), we have moreover for every $x \in N$ that

$$\lambda \omega(x^*x) = \lambda \langle T \Lambda(x), \Lambda(x) \rangle \geq \lambda_1 \langle T_1 \Lambda(x), \Lambda(x) \rangle = \lambda_1 \omega_1(x^*x)$$

so we see that $\lambda_1 \omega_1 \leq \lambda \omega$. We get in a similar way that $\lambda_2 \omega_2 \leq \lambda \omega$ ■

This allows us to give the following definition :

Definition 3.6 We define the weight φ on A such that $\varphi(x) = \sup \{ \omega(x) \mid \omega \in \mathcal{F} \}$ for every $x \in A^+$. Then φ is a densely defined lower semi-continuous weight on A such that $N \subseteq \mathcal{N}_\varphi$ and $\varphi(x^*x) \leq \|\Lambda(x)\|^2$ for every $x \in N$.

In the last proposition of this section, we will improve the relationship between φ and the mapping Λ (under some extra conditions).

Lemma 3.7 Let E be a normed space, H a Hilbert space and Λ linear mapping from within E into H . Let $(x_i)_{i \in I}$ be a net in $D(\Lambda)$ and x an element in E such that $(x_i)_{i \in I}$ converges to x and $(\Lambda(x_i))_{i \in I}$ is bounded. Then there exists a sequence $(y_n)_{n=1}^\infty$ in the convex hull of $\{x_i \mid i \in I\}$ and an element $v \in H$ such that $(y_n)_{n=1}^\infty$ converges to x and $(\Lambda(y_n))_{n=1}^\infty$ converges to v .

Proof : By the Banach-Alaoglu theorem, there exists a subnet $(x_{i_j})_{j \in J}$ of $(x_i)_{i \in I}$ and $v \in H$ such that $(\Lambda(x_{i_j}))_{j \in J}$ converges to v in the weak topology on H . (For this, we need H to be a Hilbert space.)

Fix $n \in \mathbb{N}$. Then there exists $j_n \in J$ such that $\|x_{i_{j_n}} - x\| \leq \frac{1}{n}$ for all $j \in J$ with $j \geq j_n$.

Now v belongs to the weak-closed convex hull of the set $\{\Lambda(x_{i_j}) \mid j \in J \text{ such that } j \geq j_n\}$, which is the same as the norm-closed convex hull.

Therefore, there exist $\lambda_1, \dots, \lambda_m \in \mathbb{R}^+$ with $\sum_{k=1}^m \lambda_k = 1$ and elements $\alpha_1, \dots, \alpha_m \in J$ with $\alpha_1, \dots, \alpha_m \geq j_n$ such that

$$\|v - \sum_{k=1}^m \lambda_k \Lambda(x_{i_{\alpha_k}})\| \leq \frac{1}{n}.$$

Put $y_n = \sum_{k=1}^m \lambda_k x_{i_{\alpha_k}}$. Then $y_n \in D(\Lambda)$, and $\Lambda(y_n) = \sum_{k=1}^m \lambda_k \Lambda(x_{i_{\alpha_k}})$.

Therefore, we have immediately that $\|v - \Lambda(y_n)\| \leq \frac{1}{n}$.

Furthermore,

$$\|x - y_n\| = \left\| \sum_{k=1}^m \lambda_k (x - x_{i_{\alpha_k}}) \right\| \leq \sum_{k=1}^m \lambda_k \frac{1}{n} = \frac{1}{n}$$

Therefore, we find that $(y_n)_{n=1}^\infty$ converges to x and that $(\Lambda(y_n))_{n=1}^\infty$ converges to v . ■

Using this lemma, we can prove the result we need later on.

Proposition 3.8 Suppose that

- The mapping Λ is closed.
- There exist a net $(u_i)_{i \in I}$ in N and a net $(S_i)_{i \in I}$ in $B(H)$ such that
 1. We have for every $a \in N$ and $i \in I$ that $\Lambda(au_i) = S_i \Lambda(a)$.
 2. We have for every $i \in I$ that $\|S_i\| \leq 1$.
 3. $(u_i)_{i \in I}$ converges strictly to 1 and $(S_i)_{i \in I}$ converges strongly to 1.

Then (H, Λ, π) is a GNS-construction for φ and $\mathcal{F} = \mathcal{F}_\varphi$.

We have moreover the following results :

Define for every $i \in I$ the element $\omega_i \in A_+^*$ such that $\omega_i(x) = \varphi(u_i^* x u_i)$ for $x \in A$. Then we have for every $i \in I$ that ω_i belongs to $\mathcal{F}$. Furthermore,

1. Consider $x \in \mathcal{M}_\varphi$, then $(\omega_i(x))_{i \in I}$ converges to $\varphi(x)$.

2. Consider $x \in A^+$, then $\varphi(x) = \sup \{ \omega_i(x) \mid i \in I \}$.

Proof : Choose $i \in I$. Then we have for every $a \in N$ that

$$\omega_i(a^*a) = \varphi(u_i^* a^* a u_i) = \langle \Lambda(a u_i), \Lambda(a u_i) \rangle = \langle S_i \Lambda(a), S_i \Lambda(a) \rangle . \quad (*)$$

Because $\|S_i\| \leq 1$, this implies that $\omega_i(a^*a) \leq \|\Lambda(a)\|^2$ for every $a \in N$. Therefore ω_i belongs to $\mathcal{F}$.

- Choose $a \in N$. Because $(S_i)_{i \in I}$ converges strongly to 1, equality (*) implies that $(\omega_i(a^*a))_{i \in I}$ converges to $\|\Lambda(a)\|^2$ which implies that $\|\Lambda(a)\|^2 \leq \varphi(a^*a)$, so $\varphi(a^*a) = \|\Lambda(a)\|^2$.
- Consider $a \in A$ such that the net $(\omega_i(a^*a))_{i \in I}$ is bounded.

We have for every $i \in I$ that $\omega_i \in \mathcal{F}$ which by the definition of φ implies that

$$\|\Lambda(a u_i)\|^2 = \langle \Lambda(a u_i), \Lambda(a u_i) \rangle = \varphi(u_i^* a^* a u_i) = \omega_i(a^*a)$$

Therefore the net $(\Lambda(a u_i))_{i \in I}$ is bounded.

Because $(a u_i)_{i \in I}$ converges to a , the previous lemma implies the existence of a sequence $(b_n)_{n=1}^\infty$ in $D(\Lambda)$ and $v \in H$ such that $(b_n)_{n=1}^\infty$ converges to a and $(\Lambda(b_n))_{n=1}^\infty$ converges to v . Therefore the closedness of Λ implies that a belongs to N .

Because of the definition of φ , this last result implies also that $\mathcal{N}_\varphi \subseteq N$ which implies that $N = \mathcal{N}_\varphi$. ■

4 Relatively invariant one-parameter groups

In this paper, we will have to use one-parameter groups which are relatively invariant with respect to some closed mappings arising from the GNS-construction for weights. In this section, we will gather some results about such objects.

In this section, we consider two Banach spaces E, F , a subspace N of E and a closed linear map Λ from N into F . Let α be a strongly continuous one-parameter representation on E such that there exists a strongly continuous one-parameter representation u on F together with a strictly positive number λ such that we have for every $t \in \mathbb{R}$ and $a \in N$ that $\alpha_t(a)$ belongs to N and $\Lambda(\alpha_t(a)) = \lambda^{\frac{t}{2}} u_t \Lambda(a)$.

The techniques used in the proof of the following lemma will be frequently used.

Lemma 4.1 Consider $y, z \in \mathbb{C}$, $n \in \mathbb{N}$ and $a \in N$. Then

$$\int \exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a) dt$$

belongs to N and

$$\Lambda \left(\int \exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a) dt \right) = \int \exp(-n^2(t-z)^2) \lambda^{(y+\frac{1}{2})t} u_t \Lambda(a) dt .$$

Proof : By assumption, we have for every $t \in \mathbb{R}$ that $\exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a)$ belongs to N and

$$\Lambda(\exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a)) = \exp(-n^2(t-z)^2) \lambda^{(y+\frac{1}{2})t} u_t \Lambda(a) .$$

This implies that the function

$$\mathbb{R} \rightarrow F : t \mapsto \Lambda(\exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a))$$

is integrable. Using lemma 1.1, we get that

$$\int \exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a) dt$$

belongs to N and

$$\begin{aligned} \Lambda\left(\int \exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a) dt\right) &= \int \Lambda(\exp(-n^2(t-z)^2) \lambda^{yt} \alpha_t(a)) dt \\ &= \int \exp(-n^2(t-z)^2) \lambda^{(y+\frac{1}{2})t} u_t \Lambda(a) dt. \end{aligned}$$

■

The following lemma follows easily from the previous one.

Lemma 4.2 Consider $a \in N$, $n \in \mathbb{N}$ and define

$$b = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) dt .$$

Then b is analytic with respect to α and we have for every $z \in \mathbb{C}$ that $\alpha_z(b)$ belongs to N

A first application can be found in the following result :

Proposition 4.3 Define the set

$$C = \{ a \in N \mid a \text{ is analytic with respect to } \alpha \text{ and } \alpha_z(a) \text{ belongs to } N \text{ for every } z \in \mathbb{C} \} .$$

Then

1. C is a core for Λ .
2. Let z be a complex number. If N is dense in A , then C is a core for α_z .

Proof : Define for every $n \in \mathbb{N}$ and $a \in N$ the element

$$a(n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) dt ,$$

which belongs to C by the previous lemma.

1. Choose $x \in N$. By lemma 4.1, it follows that

$$\Lambda(x(n)) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \lambda^{\frac{t}{2}} u_t \Lambda(x) dt$$

for every $n \in N$. This implies that $(\Lambda(x(n)))_{n=1}^{\infty}$ converges to $\Lambda(x)$. It is also clear that $(x(n))_{n=1}^{\infty}$ converges to x .

2. If N is dense in A , proposition 1.15 learns that the set $\langle a(n) \mid a \in N, n \in \mathbb{N} \rangle$ is a core for $D(\alpha_z)$. This implies immediately that C is a core for $D(\alpha_z)$ in this case. ■

Proposition 4.4 *Consider $z \in \mathbb{C}$ and $a \in N \cap D(\alpha_z)$ such that $\alpha_z(a)$ belongs to N . Then $\Lambda(a)$ belongs to $D(u_z)$ and $u_z \Lambda(a) = \lambda^{-\frac{z}{2}} \Lambda(\alpha_z(a))$.*

Proof : Choose $n \in \mathbb{N}$ and define

$$v_n = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) u_t \Lambda(a) dt \in F .$$

It is clear that v_n belongs to $D(u_z)$ and

$$u_z(v_n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) u_t \Lambda(a) dt .$$

By lemma 4.1, we know that

$$\frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \lambda^{-\frac{t}{2}} \alpha_t(a) dt$$

belongs to N and

$$\Lambda\left(\frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \lambda^{-\frac{t}{2}} \alpha_t(a) dt\right) = u_z(v_n) \quad (*)$$

Because a belongs to $D(\alpha_z)$, we have that

$$\frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \lambda^{-\frac{t}{2}} \alpha_t(a) dt = \lambda^{-\frac{z}{2}} \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \lambda^{-\frac{t}{2}} \alpha_t(\alpha_z(a)) dt .$$

Using lemma 4.1 once more, this implies that

$$\Lambda\left(\frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \lambda^{-\frac{t}{2}} \alpha_t(a) dt\right) = \lambda^{-\frac{z}{2}} \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) u_t \Lambda(\alpha_z(a)) dt .$$

Comparing this equality with equality (*), we get that

$$u_z(v_n) = \lambda^{-\frac{z}{2}} \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) u_t \Lambda(\alpha_z(a)) dt .$$

This implies that $(u_z(v_n))_{n=1}^{\infty}$ converges to $\lambda^{-\frac{z}{2}} \Lambda(\alpha_z(a))$. It is also clear that $(v_n)_{n=1}^{\infty}$ converges to $\Lambda(a)$. Therefore, the closedness of u_z implies that $\Lambda(a)$ belongs to $D(u_z)$ and $u_z \Lambda(a) = \lambda^{-\frac{z}{2}} \Lambda(\alpha_z(a))$. ■

Proposition 4.5 *Consider $z \in \mathbb{C}$ and $a \in N \cap D(\alpha_z)$ such that $\Lambda(a)$ belongs to $D(u_z)$. Then $\alpha_z(a)$ belongs to N and $\Lambda(\alpha_z(a)) = \lambda^{\frac{z}{2}} u_z \Lambda(a)$.*

Proof : Choose $n \in \mathbb{N}$ and define

$$b_n = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(\alpha_z(a)) dt .$$

We also have that

$$b_n = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \alpha_t(a) dt .$$

Therefore, lemma 4.1 implies that b_n belongs to N and

$$\Lambda(b_n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t-z)^2) \lambda^{\frac{t}{2}} u_t \Lambda(a) dt .$$

Because $\Lambda(a)$ belongs to $D(u_z)$, we get from the previous equation that

$$\Lambda(b_n) = \lambda^{\frac{z}{2}} \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \lambda^{\frac{t}{2}} u_t (u_z \Lambda(a)) dt .$$

It is clear from this last equation that $(\Lambda(b_n))_{n=1}^\infty$ converges to $\lambda^{\frac{z}{2}} u_z \Lambda(a)$. It is also clear from the definition that $(b_n)_{n=1}^\infty$ converges to $\alpha_z(a)$. Therefore, the closedness of Λ implies that $\alpha_z(a)$ belongs to N and $\Lambda(\alpha_z(a)) = \lambda^{\frac{z}{2}} u_z \Lambda(a)$. $\blacksquare$

Corollary 4.6 *Consider $z \in \mathbb{C}$ and $a \in N \cap D(\alpha_z)$. Then $\Lambda(a)$ belongs to $D(u_z) \Leftrightarrow \alpha_z(a)$ belongs to N .*

5 KMS-weights arising from certain GNS-constructions

In [22] and [15], Jan Verding introduced a construction procedure for weights starting from a GNS-construction. We want to repeat this procedure, but we want to end up with a KMS-weight. In order to do so, we have to impose stronger conditions on the ingredients of the construction procedure. In a last part, we describe the natural case where a KMS-weight is obtained from a left Hilbert-algebra.

First we prove some technical results.

Consider a C^* -algebra A , a Hilbert space H , a dense left ideal N of A and a closed linear mapping Λ from N into H . Let α be a norm continuous one-parameter group on A and suppose there exists a positive injective operator ∇ in H and a strictly positive number λ such that $\alpha_t(a) \in N$ and $\Lambda(\alpha_t(a)) = \lambda^{\frac{t}{2}} \nabla^{it} \Lambda(a)$ for every $t \in \mathbb{R}$ and $a \in N$.

Lemma 5.1 *Consider $a \in N \cap N^*$ and $n \in \mathbb{N}$ and define*

$$b = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) dt .$$

Then b is analytic with respect to α and $\alpha_z(b)$ belongs to $N \cap N^*$ for every $z \in \mathbb{C}$.

Proof : Choose $z \in \mathbb{C}$. By lemma 4.2, we know already that $\alpha_z(b)$ belongs to N . We also have that a^* belongs to N and that

$$b^* = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a^*) dt ,$$

so lemma 4.2 implies in this case that $\alpha_{\bar{z}}(b^*)$ belongs to N . Because $\alpha_z(b)^* = \alpha_{\bar{z}}(b^*)$, we see that $\alpha_z(b)$ belongs to N^* . $\blacksquare$

Define the set

$$C = \{ a \in N \cap N^* \mid a \text{ is analytic with respect to } \alpha \text{ and } \alpha_z(a) \text{ belongs to } N \cap N^* \text{ for every } z \in \mathbb{C} \} .$$

Proposition 5.2 *We have the following properties :*

1. *For every $x \in N \cap N^*$, there exists a sequence $(x_n)_{n=1}^\infty$ in C such that $(x_n)_{n=1}^\infty$ converges to x , $(\Lambda(x_n))_{n=1}^\infty$ converges to $\Lambda(x)$ and $(\Lambda(x_n^*))_{n=1}^\infty$ converges to $\Lambda(x^*)$.*
2. *If $N \cap N^*$ is a core for Λ , then C is a core for Λ .*
3. *Let z be a complex number. Then C is a core for α_z .*

Proof : Define for every $n \in \mathbb{N}$ and $a \in N \cap N^*$ the element

$$a(n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \alpha_t(a) dt ,$$

which belongs to C by the previous lemma.

1. Choose $x \in N \cap N^*$. Again, we see immediately that $(x(n))_{n=1}^\infty$ converges to x . Because x belongs to N , the proof of proposition 4.3 learns us that $(\Lambda(x(n)))_{n=1}^\infty$ converges to $\Lambda(x)$.
We have for every $n \in \mathbb{N}$ that $x(n)^* = x^*(n)$, so the same proof tells us that $(\Lambda(x(n)^*))_{n=1}^\infty$ converges to $\Lambda(x^*)$.
2. This follows immediately from the previous statement.
3. This is proven in the same way as the second statement of proposition 4.3.

■

The proof of the following result is due to A. Van Daele and J. Verding.

Proposition 5.3 *Consider $z \in \mathbb{C}$. Then there exists a net $(u_k)_{k \in K}$ in $C \cap A^+$ consisting of analytic elements for α and such that*

- *We have for every $k \in K$ that $\|u_k\| \leq 1$ and $\|\alpha_z(u_k)\| \leq 1$.*
- *The nets $(u_k)_{k \in K}$ and $(\alpha_z(u_k))_{k \in K}$ converge strictly to 1.*

Proof : Because N is a dense left ideal in A , we know that there exists a net $(e_q)_{q \in Q}$ in $N \cap A^+$ such that $\|e_q\| \leq 1$ for every $q \in Q$ and such that $(e_q)_{q \in Q}$ converges strictly to 1.

For every $q \in Q$ and $\delta > 0$, we define the element

$$e_q(\delta) = \frac{\delta}{\sqrt{\pi}} \int \exp(-\delta^2 t^2) \alpha_t(e_q) dt \in A^+$$

which is clearly analytic with respect to α and satisfies $\|e_q(\delta)\| \leq 1$ and $\|\alpha_z(e_q(\delta))\| \leq \exp(\delta^2 (\text{Im } z)^2)$. Lemma 5.1 implies that $e_q(\delta) \in C$ for every $q \in Q$ and $\delta > 0$.

By proposition 1.38, we have for every $\delta > 0$ that $(e_q(\delta))_{q \in Q}$ and $(\alpha_z(e_q(\delta)))_{q \in Q}$ are bounded nets which converge strictly to 1.

Let us now define the set

$$K = \{ (F, n) \mid F \text{ is a finite subset of } A \text{ and } n \in \mathbb{N} \}.$$

On K we put an order such that

$$(F_1, n_1) \leq (F_2, n_2) \Leftrightarrow F_1 \subseteq F_2 \text{ and } n_1 \leq n_2$$

for every $(F_1, n_1), (F_2, n_2) \in K$. In this way, K becomes a directed set.

Let us fix $k = (F, n) \in K$.

Firstly, there exist an element $\delta_k > 0$ such that $|\exp(-\delta_k^2 (\operatorname{Im} z)^2) - 1| \leq \frac{1}{2n} \frac{1}{\|x\|+1}$ for every $x \in F$.

Secondly, there exist $q_k \in Q$ such that

- We have that $\|e_{q_k}(\delta_k) x - x\| \leq \frac{1}{2n}$ for every $x \in F$.
- We have that $\|\alpha_z(e_{q_k}(\delta_k)) x - x\| \leq \frac{1}{2n}$ for every $x \in F$.
- We have that $\|x \alpha_z(e_{q_k}(\delta_k)) - x\| \leq \frac{1}{2n}$ for every $x \in F$.

Now we define $u_k = \exp(-\delta_k^2 (\operatorname{Im} z)^2) e_{q_k}(\delta_k) \in C \cap A^+$.

It follows that u_k is analytic with respect to α and $\|u_k\| \leq 1$. We have also that

$$\begin{aligned} \|\alpha_z(u_k)\| &= \exp(-\delta_k^2 (\operatorname{Im} z)^2) \|\alpha_z(e_{q_k}(\delta_k))\| \\ &\leq \exp(-\delta_k^2 (\operatorname{Im} z)^2) \exp(-\delta_k^2 (\operatorname{Im} z)^2) = 1 \end{aligned}$$

Now we prove that $(u_k)_{k \in K}$ converges strictly to 1.

Choose $x \in A$ and $\varepsilon > 0$. Then there exists n_0 in $\mathbb{N}$ such that $\frac{1}{n_0} \leq \varepsilon$. Put $k_0 = (\{x\}, n_0) \in K$.

Take $k = (F, n) \in K$ such that $k \geq k_0$, so x belongs to F and $n_0 \leq n$. Therefore,

$$\begin{aligned} \|u_k x - x\| &= \|\exp(-\delta_k^2 (\operatorname{Im} z)^2) e_{q_k}(\delta_k) x - x\| \\ &\leq \|\exp(-\delta_k^2 (\operatorname{Im} z)^2) e_{q_k}(\delta_k) x - \exp(-\delta_k^2 (\operatorname{Im} z)^2) x\| + \|\exp(-\delta_k^2 (\operatorname{Im} z)^2) x - x\| \\ &= \exp(-\delta_k^2 (\operatorname{Im} z)^2) \|e_{q_k}(\delta_k) x - x\| + |\exp(-\delta_k^2 (\operatorname{Im} z)^2) - 1| \|x\| \\ &\leq \|e_{q_k}(\delta_k) x - x\| + \frac{1}{2n(\|x\|+1)} \|x\| \leq \frac{1}{2n} + \frac{1}{2n} = \frac{1}{n} \leq \varepsilon. \end{aligned}$$

Because $(u_k)_{k \in K}$ consists of selfadjoint elements, this implies that $(u_k)_{k \in K}$ converges strictly to 1.

Completely analogously, one proves that $(\alpha_z(u_k) x)_{k \in K}$ converges to x for every $x \in A$ (just replace u_k by $\alpha_z(u_k)$ in the proof above). Similarly one proves that $(x \alpha_z(u_k))_{k \in K}$ converges to x for every $x \in A$. Consequently, we have that $(\alpha_z(u_k))_{k \in K}$ converges strictly to 1. $\blacksquare$

5.1 The first construction procedure.

In the first construction procedure, we will use the following ingredients:

Consider a C^* -algebra A and a Hilbert space H . Let N be a dense left ideal of A , Λ a closed linear mapping from N into H with dense range and π a non-degenerate representation of A on H such that $\pi(x)\Lambda(a) = \Lambda(xa)$ for every $x \in A$ and $a \in N$.

Furthermore, assume the existence of

- a norm continuous one-parameter group σ on A
- an injective positive operator ∇ in H

such that:

- We have for every $a \in N$ and $t \in \mathbb{R}$ that $\sigma_t(a)$ belongs to N and $\Lambda(\sigma_t(a)) = \nabla^{it} \Lambda(a)$.
- There exists a core K for Λ such that

1. $K \subseteq D(\sigma_{\frac{i}{2}})$, $\sigma_{\frac{i}{2}}(K)^* \subseteq N$ and $\|\Lambda(x)\| = \|\Lambda(\sigma_{\frac{i}{2}}(x)^*)\|$ for every $x \in K$.
2. $\sigma_t(K) \subseteq K$ for every $t \in \mathbb{R}$.

Our first objective is to extend the property about K to $N \cap D(\sigma_{\frac{i}{2}})$.

In this section, we will use the following notation: For every $a \in A$ and $n \in \mathbb{N}$, we put

$$a(n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \sigma_t(a) dt ,$$

it is clear that $a(n)$ is analytic with respect to σ .

Furthermore, we know from lemma 4.1 that we have for every $a \in N$ and $n \in \mathbb{N}$ that $a(n)$ belongs to N and

$$\Lambda(a(n)) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \nabla^{it} \Lambda(a) dt. \quad (1)$$

Lemma 5.4 *We have the following properties:*

1. We have for every $x \in K$ and $n \in \mathbb{N}$ that $x(n) \in N \cap D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(x(n))^* \in N$.
2. We have for every $x, y \in K$ and $m, n \in \mathbb{N}$ that

$$\langle \Lambda(y(n)), \Lambda(x(m)) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(x(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y(n))^*) \rangle .$$

Proof :

1. Choose $x \in K$ and $n \in \mathbb{N}$. By equation 1, we know already that $x(n)$ belongs to N . We also have that $x(n)$ belongs to $D(\sigma_{\frac{i}{2}})$. The fact that x belongs to $D(\sigma_{\frac{i}{2}})$ implies that

$$\sigma_{\frac{i}{2}}(x(n))^* = (\sigma_{\frac{i}{2}}(x)(n))^* = \sigma_{\frac{i}{2}}(x)^*(n) \quad (a)$$

Therefore, equation 1 and the fact that $\sigma_{\frac{i}{2}}(x)^* \in N$ imply that $\sigma_{\frac{i}{2}}(x(n))^*$ belongs to N .

2. Choose $x, y \in K$ and $m, n \in \mathbb{N}$. Using equation (a) from the first part of the proof and equation 1 once more, we get the following equalities

$$\begin{aligned} \langle \Lambda(\sigma_{\frac{i}{2}}(x(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y(n))^*) \rangle &= \langle \Lambda(\sigma_{\frac{i}{2}}(x)^*(m)), \Lambda(\sigma_{\frac{i}{2}}(y)^*(n)) \rangle \\ &= \frac{m n}{\pi} \int \int \exp(-(m^2 s^2 + n^2 t^2)) \langle \nabla^{is} \Lambda(\sigma_{\frac{i}{2}}(x)^*), \nabla^{it} \Lambda(\sigma_{\frac{i}{2}}(y)^*) \rangle ds dt \end{aligned} \quad (b)$$

By polarisation, we get that

$$\langle \Lambda(a), \Lambda(b) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(b)^*), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle$$

for every $a, b \in K$.

Fix $s, t \in \mathbb{R}$. By assumption, we know that $\sigma_s(x), \sigma_t(y)$ belong to K . So, using the previous equality, we see that

$$\begin{aligned} \langle \nabla^{is} \Lambda(\sigma_{\frac{i}{2}}(x)^*), \nabla^{it} \Lambda(\sigma_{\frac{i}{2}}(y)^*) \rangle &= \langle \Lambda(\sigma_s(\sigma_{\frac{i}{2}}(x)^*)), \Lambda(\sigma_t(\sigma_{\frac{i}{2}}(y)^*)) \rangle \\ &= \langle \Lambda(\sigma_{\frac{i}{2}}(\sigma_s(x))^*), \Lambda(\sigma_{\frac{i}{2}}(\sigma_t(y))^*) \rangle = \langle \Lambda(\sigma_t(y)), \Lambda(\sigma_s(x)) \rangle = \langle \nabla^{it} \Lambda(y), \nabla^{is} \Lambda(x) \rangle \end{aligned}$$

Substituting this equality into equation (b) gives us that

$$\begin{aligned} \langle \Lambda(\sigma_{\frac{i}{2}}(x(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y(n))^*) \rangle &= \frac{m n}{\pi} \int \int \exp(-(m^2 s^2 + n^2 t^2)) \langle \nabla^{it} \Lambda(y), \nabla^{is} \Lambda(x) \rangle ds dt \\ &= \langle \Lambda(y(n)), \Lambda(x(m)) \rangle , \end{aligned}$$

where we used equation 1 in the last equality.

■

Lemma 5.5 *We have the following properties:*

1. We have for every $x \in N$ and $n \in \mathbb{N}$ that $x(n) \in N \cap D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(x(n))^* \in N$.
2. We have for every $x, y \in N$ and $m, n \in \mathbb{N}$ that

$$\langle \Lambda(y(n)), \Lambda(x(m)) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(x(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y(n))^*) \rangle .$$

Proof :

1. Choose $x \in N$ and $n \in \mathbb{N}$. We know by equation 1 that $x(n)$ belongs to N . We have also that $x(n)$ belongs to $D(\sigma_{\frac{i}{2}})$.

Because K is a core for Λ , there exists a sequence $(x_k)_{k=1}^\infty$ in K such that $(x_k)_{k=1}^\infty \rightarrow x$ and $(\Lambda(x_k))_{k=1}^\infty \rightarrow \Lambda(x)$. From the previous lemma we know that $x_k(n)$ belongs to $N \cap D(\sigma_{\frac{i}{2}})$ and that $\sigma_{\frac{i}{2}}(x_k(n))^*$ belongs to N for every $k \in \mathbb{N}$.

For every $k \in \mathbb{N}$, we have that

$$\sigma_{\frac{i}{2}}(x_k(n))^* = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t + \frac{i}{2})^2) \sigma_t(x_k^*) dt$$

and similarly,

$$\sigma_{\frac{i}{2}}(x(n))^* = \frac{n}{\sqrt{\pi}} \int \exp(-n^2(t + \frac{i}{2})^2) \sigma_t(x^*) dt .$$

Comparing these two equations and remembering that $(x_k)_{k=1}^\infty$ converges to x , we get that $(\sigma_{\frac{i}{2}}(x_k(n))^*)_{k=1}^\infty$ converges to $\sigma_{\frac{i}{2}}(x(n))^*$. (a)

Because $(\Lambda(x_k))_{k=1}^\infty$ converges to $\Lambda(x)$, it is clear from equation 1 that $(\Lambda(x_k(n)))_{k=1}^\infty$ converges to $\Lambda(x(n))$. By the previous lemma, we have for every $k, l \in \mathbb{N}$ that

$$\|\Lambda(\sigma_{\frac{i}{2}}(x_l(n))^*) - \Lambda(\sigma_{\frac{i}{2}}(x_k(n))^*)\| = \|\Lambda(x_l(n)) - \Lambda(x_k(n))\|$$

Therefore, the convergence of $(\Lambda(x_k(n)))_{k=1}^\infty$ implies that $(\Lambda(\sigma_{\frac{i}{2}}(x_k(n))^*))_{k=1}^\infty$ is a Cauchy sequence and hence convergent. (b)

Using the closedness of Λ and conclusions (a) and (b), we get that $\sigma_{\frac{i}{2}}(x(n))^*$ belongs to N and that $(\Lambda(\sigma_{\frac{i}{2}}(x_k(n))^*))_{k=1}^\infty$ converges to $\Lambda(\sigma_{\frac{i}{2}}(x(n))^*)$.

2. Choose $x, y \in N$ and $m, n \in \mathbb{N}$. Then there exists sequences $(x_k)_{k=1}^\infty, (y_k)_{k=1}^\infty$ in K such that $(x_k)_{k=1}^\infty \rightarrow x, (y_k)_{k=1}^\infty \rightarrow y, (\Lambda(x_k))_{k=1}^\infty \rightarrow \Lambda(x)$ and $(\Lambda(y_k))_{k=1}^\infty \rightarrow \Lambda(y)$.

By the first part of the proof, we know that $(\Lambda(x_k(m)))_{k=1}^\infty \rightarrow \Lambda(x(m)), (\Lambda(y_k(n)))_{k=1}^\infty \rightarrow \Lambda(y(n)), (\Lambda(\sigma_{\frac{i}{2}}(x_k(m))^*))_{k=1}^\infty \rightarrow \Lambda(\sigma_{\frac{i}{2}}(x(m))^*)$ and $(\Lambda(\sigma_{\frac{i}{2}}(y_k(n))^*))_{k=1}^\infty \rightarrow \Lambda(\sigma_{\frac{i}{2}}(y(n))^*)$. By the previous lemma, we know that

$$\langle \Lambda(y_k(n)), \Lambda(x_k(m)) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(x_k(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y_k(n))^*) \rangle$$

for every $k \in \mathbb{N}$. This implies that

$$\langle \Lambda(y(n)), \Lambda(x(m)) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(x(m))^*), \Lambda(\sigma_{\frac{i}{2}}(y(n))^*) \rangle .$$

■

At last, we arrive at a final form.

Proposition 5.6 *Consider $x \in N \cap D(\sigma_{\frac{i}{2}})$. Then $\sigma_{\frac{i}{2}}(x)^*$ belongs to N and $\|\Lambda(x)\| = \|\Lambda(\sigma_{\frac{i}{2}}(x)^*)\|$.*

Proof : By the previous lemma, we have for every $n \in \mathbb{N}$ that $x(n)$ belongs to $N \cap D(\sigma_{\frac{i}{2}})$, $\sigma_{\frac{i}{2}}(x(n))^*$ belongs to N and $\|\Lambda(x(n))\| = \|\Lambda(\sigma_{\frac{i}{2}}(x(n))^*)\|$.

By equation 1, we have for every $n \in \mathbb{N}$ that

$$\Lambda(x(n)) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \nabla^{it} \Lambda(x) dt$$

which implies that $(\Lambda(x(n)))_{n=1}^{\infty} \rightarrow \Lambda(x)$.

By the previous lemma, we have for every $m, n \in \mathbb{N}$ that

$$\|\Lambda(\sigma_{\frac{i}{2}}(x(m))^*) - \Lambda(\sigma_{\frac{i}{2}}(x(n))^*)\| = \|\Lambda(x(m)) - \Lambda(x(n))\|.$$

Hence, the convergence of $(\Lambda(x(n)))_{n=1}^{\infty}$ implies that $(\Lambda(\sigma_{\frac{i}{2}}(x(n))^*))_{n=1}^{\infty}$ is a Cauchy sequence and hence convergent.

Because x belongs to $D(\sigma_{\frac{i}{2}})$, we have for every $n \in \mathbb{N}$ that

$$\sigma_{\frac{i}{2}}(x(n))^* = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \sigma_t(\sigma_{\frac{i}{2}}(x)^*) dt,$$

from which it follows easily that $(\sigma_{\frac{i}{2}}(x(n))^*)_{n=1}^{\infty}$ converges to $\sigma_{\frac{i}{2}}(x)^*$.

Hence, the closedness of Λ implies that $\sigma_{\frac{i}{2}}(x)^*$ belongs to N and $(\Lambda(\sigma_{\frac{i}{2}}(x(n))^*))_{n=1}^{\infty} \rightarrow \Lambda(\sigma_{\frac{i}{2}}(x)^*)$.

Because $\|\Lambda(\sigma_{\frac{i}{2}}(x(n))^*)\| = \|\Lambda(x(n))\|$ for every $n \in \mathbb{N}$, we will also have that $\|\Lambda(\sigma_{\frac{i}{2}}(x)^*)\| = \|\Lambda(x)\|$. ■

Using the fact that $\sigma_{\frac{i}{2}}(\sigma_{\frac{i}{2}}(x)^*) = x$ for every $x \in D(\sigma_{\frac{i}{2}})$, the following corollary follows readily.

Corollary 5.7 *Consider $x \in D(\sigma_{\frac{i}{2}})$. Then x belongs to N if and only if $\sigma_{\frac{i}{2}}(x)^*$ belongs to N .*

Because $K \subseteq N \cap D(\sigma_{\frac{i}{2}})$, we have that $N \cap D(\sigma_{\frac{i}{2}})$ is a core for Λ . Therefore, $\Lambda(N \cap D(\sigma_{\frac{i}{2}}))$ will certainly be dense in H . This allows us to introduce the following definition.

Definition 5.8 *We define the anti-unitary operator J on H such that $J\Lambda(x) = \Lambda(\sigma_{\frac{i}{2}}(x)^*)$ for every $x \in N \cap D(\sigma_{\frac{i}{2}})$.*

It follows immediately that $J\Lambda(N \cap D(\sigma_{\frac{i}{2}})) \subseteq \Lambda(N \cap D(\sigma_{\frac{i}{2}}))$ and that $J(J\Lambda(x)) = \Lambda(x)$ for every $x \in N \cap D(\sigma_{\frac{i}{2}})$. This implies that $J^2 = 1$.

The next proposition is crucial in the construction of a weight out of our ingredients.

Proposition 5.9 *Consider $a \in D(\sigma_{\frac{i}{2}})$ and $x \in N$. Then xa belongs to N and $\Lambda(xa) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x)$*

Proof : Choose $b, y \in N \cap D(\sigma_{\frac{i}{2}})$. Then we have that $yb \in N \cap D(\sigma_{\frac{i}{2}})$. So, by the definition of J , we have that

$$\begin{aligned} J\Lambda(yb) &= \Lambda(\sigma_{\frac{i}{2}}(yb)^*) = \Lambda(\sigma_{\frac{i}{2}}(b)^* \sigma_{\frac{i}{2}}(y)^*) \\ &= \pi(\sigma_{\frac{i}{2}}(b)^*) \Lambda(\sigma_{\frac{i}{2}}(y)^*) = \pi(\sigma_{\frac{i}{2}}(b))^* J\Lambda(y). \end{aligned}$$

Therefore, $\Lambda(yb) = J\pi(\sigma_{\frac{i}{2}}(b))^* J\Lambda(y)$.

Proposition 4.3 implies the existence of sequences $(a_n)_{n=1}^\infty$, $(x_n)_{n=1}^\infty$ in $N \cap D(\sigma_{\frac{i}{2}})$ such that $(a_n)_{n=1}^\infty$ converges to a , $(x_n)_{n=1}^\infty$ converges to x , $(\sigma_{\frac{i}{2}}(a_n))_{n=1}^\infty$ converges to $\sigma_{\frac{i}{2}}(a)$ and $(\Lambda(x_n))_{n=1}^\infty$ converges to $\Lambda(x)$.

This implies immediately that $(x_n a_n)_{n=1}^\infty$ converges to xa .

By the first part of this proof, we know that $x_n a_n$ belongs to N and

$$\Lambda(x_n a_n) = J\pi(\sigma_{\frac{i}{2}}(a_n))^* J\Lambda(x_n)$$

for every $n \in \mathbb{N}$. This implies that

$$(\Lambda(x_n a_n))_{n=1}^\infty \rightarrow J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x) .$$

The closedness of Λ implies that xa belongs to N and $\Lambda(xa) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x)$. ■

Define the set

The following lemma is due to J. Verding and can be found in [22].

Lemma 5.10 *There exists a net $(u_k)_{k \in K} \in N \cap A^+$ and a net $(T_k)_{k \in K}$ in $\pi(A)'$ such that*

1. *We have for every $k \in K$ that $\|u_k\| \leq 1$ and $\|T_k\| \leq 1$.*
2. *We have for every $k \in k$ that u_k is analytic with respect to σ .*
3. *$(u_k)_{k \in K}$ converges strictly to 1 and $(T_k)_{k \in K}$ converges strongly* to 1.*
4. *For every $a \in N$ and $k \in K$, we have that $\Lambda(au_k) = T_k \Lambda(a)$.*

Proof : By proposition 5.3 we get the existence of a net $(u_k)_{k \in K}$ in $N \cap A^+$ consisting of elements which are analytic with respect to σ and such that

- We have for every $k \in K$ that $\|u_k\| \leq 1$ and $\|\sigma_{\frac{i}{2}}(u_k)\| \leq 1$.
- The nets $(u_k)_{k \in K}$ and $(\sigma_{\frac{i}{2}}(u_k))_{k \in K}$ converge strictly to 1.

Choose $k \in K$ and define the element $T_k = J\pi(\sigma_{\frac{i}{2}}(u_k))^* J \in B(H)$. We have immediately that $\|T_k\| \leq 1$. Proposition 5.9 implies that $\Lambda(au_k) = T_k \Lambda(a)$ for every $a \in N$.

We have also that T_k belongs to $\pi(A)'$: Choose $a \in A$. Then we have for every $b \in N$ that

$$\pi(a)T_k \Lambda(b) = \pi(a)\Lambda(bu_k) = \pi(ab)\Lambda(u_k) = T_k \Lambda(ab) = T_k \pi(a)\Lambda(b) .$$

Hence, $\pi(a)T_k = T_k \pi(a)$.

It is clear that $(T_k)_{k \in K}$ converges strongly* to 1. ■

This proposition allows us to use the construction procedure of J.Verding to get a weight (see definition 3.6 and proposition 3.8).

Definition 5.11 *Define the set*

$$\mathcal{F} = \{ \omega \in A_+^* \mid \omega(a^*a) \leq \|\Lambda(a)\|^2 \text{ for every } a \in N \} .$$

We define the mapping φ from A^+ to $[0, \infty]$ such that

$$\varphi(x) = \sup\{\omega(x) \mid \omega \in \mathcal{F}\}$$

for every $x \in A^+$. Then φ is a densely defined lower semi-continuous weight on A . We have in this case also that $\mathcal{N}_\varphi = N$ and $\varphi(b^*a) = \langle \Lambda(a), \Lambda(b) \rangle$ for every $a, b \in N$. Therefore, (H, π, Λ) is a GNS-construction for φ .

Proposition 5.6 implies immediately that φ is a KMS-weight with respect to σ .

5.2 The second construction procedure.

In the next part of this section, we want to give a second construction procedure for KMS-weights, starting with slightly different conditions on the ingredients than in the previous case.

In this section we will use the following ingredients:

Consider a C^* -algebra A and a Hilbert space H . Let N be a dense left ideal of A , Λ a closed linear mapping from N into H with dense range and π a non-degenerate representation of A on H such that $\pi(x)\Lambda(a) = \Lambda(xa)$ for every $x \in A$ and $a \in N$.

Furthermore, assume the existence of

- a norm continuous one-parameter group σ on A
- an injective positive operator ∇ in H
- an anti-unitary operator I on H

such that:

- We have for every $a \in N$ and $t \in \mathbb{R}$ that $\sigma_t(a)$ belongs to N and $\Lambda(\sigma_t(a)) = \nabla^{it}\Lambda(a)$.
- We have for every $x \in N$ and $a \in D(\sigma_{\frac{i}{2}})$ that xa belongs to N and $\Lambda(xa) = I\pi(\sigma_{\frac{i}{2}}(a))^*I^*\Lambda(x)$.

If we examine the proof of lemma 5.10 we see that we only use proposition 5.9 and the fact that Λ is invariant under σ , so the conclusion of lemma 5.10 still holds in this case. This allows us to define a weight φ in the same way as before (see definition 3.6 and proposition 3.8).

Definition 5.12 *Define the set*

$$\mathcal{F} = \{ \omega \in A_+^* \mid \omega(a^*a) \leq \|\Lambda(a)\|^2 \text{ for every } a \in N \} .$$

We define the mapping φ from A^+ to $[0, \infty]$ such that

$$\varphi(x) = \sup\{ \omega(x) \mid \omega \in \mathcal{F} \}$$

for every $x \in A^+$. Then φ is a densely defined lower semi-continuous weight on A . We have that $\mathcal{N}_\varphi = N$ and $\varphi(b^*a) = \langle \Lambda(a), \Lambda(b) \rangle$ for every $a, b \in N$. Therefore, (H, π, Λ) is a GNS-construction for φ .

We still have to prove that φ is a KMS-weight with respect to σ .

Define the set

$$C = \{ a \in N \cap N^* \mid a \text{ is analytic with respect to } \sigma \text{ and } \sigma_z(a) \text{ belongs to } N \cap N^* \text{ for every } z \in \mathbb{C} \} .$$

Then C is a sub- $*$ -algebra of A such that $C \subset N \cap D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(C)^* \subseteq N$. It is also clear that $\sigma_z(C) \subseteq C$ for every $z \in \mathbb{C}$.

Lemma 5.13 *We have that C is a core for Λ .*

Proof : We prove first that $N \cap N^*$ is a core for Λ .

Choose $x \in N$. Because N^* is dense in A , there exists a bounded net $(e_k)_{k \in K}$ in N^* such that $(e_k)_{k \in K}$ converges strictly to 1. For every $k \in K$, we have that $e_k a$ belongs to $N^* N$, which is a subset of $N \cap N^*$. Because $\Lambda(e_k a) = \pi(e_k) \Lambda(a)$ for every $k \in K$, we see that $(\Lambda(e_k a))_{k \in K}$ converges to $\Lambda(a)$. It is also clear that $(e_k a)_{k \in K}$ converges to a .

Therefore, proposition 5.2 implies that C is a core for Λ . ■

Proposition 5.14 *We have that φ is a KMS-weight with modular group σ .*

Proof : Choose $a \in C$.

Take $x \in C$. Because a belongs to $D(\sigma_{\frac{i}{2}})$, we have by assumption that

$$\begin{aligned} \langle \pi(x) \Lambda(a), \Lambda(a) \rangle &= \langle I \pi(\sigma_{\frac{i}{2}}(a))^* I^* \Lambda(x), \Lambda(a) \rangle = \langle \Lambda(x), I \pi(\sigma_{\frac{i}{2}}(a)) I^* \Lambda(a) \rangle \\ &= \langle \Lambda(x), I \pi(\sigma_{-\frac{i}{2}}(a^*))^* I^* \Lambda(a) \rangle = \langle \Lambda(x), I \pi(\sigma_{\frac{i}{2}}(\sigma_{-i}(a^*)))^* I^* \Lambda(a) \rangle \\ &= \langle \Lambda(x), \pi(a) \Lambda(\sigma_{-i}(a^*)) \rangle = \langle \Lambda(a^* x), \Lambda(\sigma_{-i}(a^*)) \rangle, \quad (*) \end{aligned}$$

where, in the second last equality, we used one of the assumptions of this second construction procedure once again. We know that $\sigma_{-\frac{i}{2}}(a^*)$ belongs to $N \cap D(\sigma_{-\frac{i}{2}})$ and $\sigma_{-\frac{i}{2}}(\sigma_{-\frac{i}{2}}(a^*)) = \sigma_{-i}(a^*) \in N$. Hence, proposition 4.4 implies that $\Lambda(\sigma_{-\frac{i}{2}}(a^*))$ belongs to $D(\nabla^{\frac{1}{2}})$ and

$$\nabla^{\frac{1}{2}} \Lambda(\sigma_{-\frac{i}{2}}(a^*)) = \Lambda(\sigma_{-i}(a^*)).$$

Similarly, we have that $a^* x$ belongs to $N \cap D(\sigma_{-\frac{i}{2}})$ and $\sigma_{-\frac{i}{2}}(a^* x)$ belongs to N . Therefore, $\Lambda(a^* x)$ belongs to $D(\nabla^{\frac{1}{2}})$ and

$$\nabla^{\frac{1}{2}} \Lambda(a^* x) = \Lambda(\sigma_{-\frac{i}{2}}(a^* x)).$$

Using these two results, equation (*) implies that

$$\begin{aligned} \langle \pi(x) \Lambda(a), \Lambda(a) \rangle &= \langle \Lambda(a^* x), \nabla^{\frac{1}{2}} \Lambda(\sigma_{-\frac{i}{2}}(a^*)) \rangle = \langle \nabla^{\frac{1}{2}} \Lambda(a^* x), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle \\ &= \langle \Lambda(\sigma_{-\frac{i}{2}}(a^* x)), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle = \langle \Lambda(\sigma_{-\frac{i}{2}}(a^*) \sigma_{-\frac{i}{2}}(x)), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle \\ &= \langle I \pi(\sigma_{\frac{i}{2}}(\sigma_{-\frac{i}{2}}(x)))^* I^* \Lambda(\sigma_{-\frac{i}{2}}(a^*)), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle \\ &= \langle I \pi(x)^* I^* \Lambda(\sigma_{\frac{i}{2}}(a)^*), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle. \end{aligned}$$

Because C is dense in A and π is non-degenerate, this equality implies easily that we can replace x in this equality by 1. Hence,

$$\langle \Lambda(a), \Lambda(a) \rangle = \langle \Lambda(\sigma_{\frac{i}{2}}(a)^*), \Lambda(\sigma_{\frac{i}{2}}(a)^*) \rangle.$$

The conclusion follows from proposition 5.6. ■

5.3 The last one.

In the last part of this section, we describe a natural way to get a KMS-weight on a C^* -algebra from a left Hilbert algebra.

Consider a Hilbert space H and a left Hilbert algebra $\mathcal{U}$ on H with modular operator ∇ , modular conjugation J and associated faithful semi-finite normal weight $\tilde{\varphi}$ on $\mathcal{L}(\mathcal{U})$. Hence

$$\mathcal{N}_{\tilde{\varphi}} = \{ L_v \mid v \in H \text{ such that } v \text{ is left bounded with respect to } \mathcal{U} \}$$

and $\tilde{\varphi}((L_w)^*(L_v)) = \langle v, w \rangle$ for every v, w in H which are left bounded with respect to $\mathcal{U}$. We have also that

$$\mathcal{N}_{\tilde{\varphi}} \cap \mathcal{N}_{\tilde{\varphi}}^* = \{ L_v \mid v \in \mathcal{U}'' \}.$$

Let A be a C^* -algebra and π a nondegenerate $*$ -representation of A on H such that $\pi(A) \subseteq \mathcal{L}(\mathcal{U})$ and suppose there exists a one-parameter group σ on A such that $\pi(\sigma_t(a)) = \nabla^{it}\pi(a)\nabla^{-it}$ for every $t \in \mathbb{R}$ and $a \in A$.

By definition, $\tilde{\varphi}\pi$ will denote the mapping $\tilde{\varphi} \circ (\pi|_{A^+})$ from A^+ into $[0, \infty]$. We put $\varphi = \tilde{\varphi}\pi$. Then φ is a lower semi-continuous weight on A such that

- $\mathcal{M}_{\varphi}^+ = \{ a \in A \mid \pi(a) \text{ belongs to } \mathcal{M}_{\tilde{\varphi}}^+ \}$
- $\mathcal{M}_{\varphi} \subseteq \{ a \in A \mid \pi(a) \text{ belongs to } \mathcal{M}_{\tilde{\varphi}} \}$
- $\mathcal{N}_{\varphi} = \{ a \in A \mid \pi(a) \text{ belongs to } \mathcal{N}_{\tilde{\varphi}} \}$
- $\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^* = \{ a \in A \mid \pi(a) \text{ belongs to } \mathcal{N}_{\tilde{\varphi}} \cap \mathcal{N}_{\tilde{\varphi}}^* \}$

and $\varphi(a) = \tilde{\varphi}(\pi(a))$ for every $a \in \mathcal{M}_{\varphi}$.

Define the following mapping Λ from $\mathcal{N}_{\varphi}$ into H : For every $a \in \mathcal{N}_{\varphi}$, there exists a unique element $v \in H$ such that v is left bounded with respect to $\mathcal{U}$ and $L_v = \pi(a)$ and we define $\Lambda(a) = v$, so we get that $L_{\Lambda(a)} = \pi(a)$.

In the following, we will suppose that $\mathcal{N}_{\varphi}$ is dense in A and that $\Lambda(\mathcal{N}_{\varphi})$ is dense in H .

It is not difficult to check in this case that (H, Λ, π) is a GNS-construction for φ .

It follows easily that $\Lambda(\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*) \subseteq \mathcal{U}''$ and $\Lambda(a)^* = \Lambda(a^*)$ for every $a \in \mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*$.

Proposition 5.15 *We have that φ is a KMS-weight with modular group σ and such that $\Lambda(\sigma_t(a)) = \nabla^{it}\Lambda(a)$ for every $a \in \mathcal{N}_{\varphi}$.*

Proof :

1. Choose $a \in \mathcal{N}_{\varphi}$ and $t \in \mathbb{R}$. Then $\Lambda(a)$ is left bounded with respect to $\mathcal{U}$ and $L_{\Lambda(a)} = \pi(a)$. So $\pi(\sigma_t(a)) = \nabla^{it}\pi(a)\nabla^{-it} = \nabla^{it}L_{\Lambda(a)}\nabla^{-it}$. Hence, the Tomita theorem implies that $\nabla^{it}\Lambda(a)$ is left bounded with respect to $\mathcal{U}$ and $L_{\nabla^{it}\Lambda(a)} = \nabla^{it}L_{\Lambda(a)}\nabla^{-it} = \pi(\sigma_t(a))$. This implies that $\sigma_t(a)$ belongs to $\mathcal{N}_{\varphi}$ and $\Lambda(\sigma_t(a)) = \nabla^{it}\Lambda(a)$.

From this, we get immediately that φ is invariant under σ .

2. Choose $b \in D(\sigma_{-\frac{i}{2}}) \cap \mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*$ and $y \in \mathcal{N}_{\varphi}^* \cap \mathcal{N}_{\varphi}$. Then y^*b belongs to $\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*$.

The function $S(-\frac{i}{2}) \rightarrow B(H) : u \mapsto \pi(\sigma_u(b))$ is continuous on $S(-\frac{i}{2})$, analytic on $S(-\frac{i}{2})^0$ and $\pi(\sigma_t(b)) = \nabla^{it}\pi(b)\nabla^{-it}$ for every $t \in \mathbb{R}$ (by assumption). This implies that $D(\nabla^{\frac{1}{2}}\pi(b)\nabla^{-\frac{1}{2}}) = D(\nabla^{-\frac{1}{2}})$ and $\nabla^{\frac{1}{2}}\pi(b)\nabla^{-\frac{1}{2}} \subseteq \pi(\sigma_{-\frac{i}{2}}(b))$. (a)

Because y belongs to $\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*$, $\Lambda(y)$ belongs to $\mathcal{U}''$ and $\Lambda(y)^* = \Lambda(y^*)$. Therefore we have that $J\Lambda(y)$ belongs to $D(\nabla^{-\frac{1}{2}})$ and $\nabla^{-\frac{1}{2}}J\Lambda(y) = \Lambda(y^*)$. Using (a), we have also that $\pi(b)\nabla^{-\frac{1}{2}}J\Lambda(y)$ belongs to $D(\nabla^{\frac{1}{2}})$ and $\nabla^{\frac{1}{2}}(\pi(b)\nabla^{-\frac{1}{2}}J\Lambda(y)) = \pi(\sigma_{-\frac{i}{2}}(b))J\Lambda(y)$.

These two results imply together that $\Lambda(by^*)$ belongs to $D(\nabla^{\frac{1}{2}})$ and $\nabla^{\frac{1}{2}}\Lambda(by^*) = \pi(\sigma_{-\frac{i}{2}}(b))J\Lambda(y)$

Because by^* belongs to $\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi}^*$, we have that $J\nabla^{\frac{1}{2}}\Lambda(by^*) = \Lambda(yb^*)$. Substituting this in the last equality gives us that

$$\Lambda(yb^*) = J\pi(\sigma_{-\frac{i}{2}}(b))J\Lambda(y) \quad (b)$$

Because $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ is dense in A , proposition 5.2 implies that $D(\sigma_{-\frac{i}{2}}) \cap \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ is a core for $\sigma_{-\frac{i}{2}}$. We know also that $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ is a core for Λ . Using these two facts and equality (b), it is now rather easy to prove for every $a \in D(\sigma_{-\frac{i}{2}})$ and $x \in \mathcal{N}_\varphi$ that xa^* belongs to $\mathcal{N}_\varphi$ and $\Lambda(xa^*) = J\pi(\sigma_{-\frac{i}{2}}(a))J\Lambda(x)$.

Proposition 5.14 implies that φ is KMS with respect to σ . ■

Proposition 5.16 *We have that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a sub left Hilbert algebra of $\mathcal{U}''$ and $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)'' = \mathcal{U}''$.*

Proof : Because φ is a densely defined lower semi-continuous weight on A , we know that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is dense in H . By a result of [18], we see that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a sub left Hilbert algebra of $\mathcal{U}''$. Choose $n \in \mathbb{N}$ and $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. Define

$$a(n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \sigma_t(a) dt .$$

Lemma 4.1 implies that $a(n)$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ and

$$\Lambda(a(n)) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \nabla^{it} \Lambda(a) dt .$$

Because $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is dense in H , this equation implies that the set $\langle \Lambda(a(n)) \mid n \in \mathbb{N}, a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \rangle$ is a core for $\nabla^{\frac{1}{2}}$. Therefore, we get that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a core for $\nabla^{\frac{1}{2}}$. Hence, left Hilbert algebra theory implies that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)'' = \mathcal{U}''$. ■

6 Properties of KMS-weights

In this section we will prove important properties of KMS-weights but we will start of with some equivalent definitions of KMS-weights.

Proposition 6.1 *Consider a C^* -algebra A and a densely defined weight φ on A with GNS-construction (H, Λ, π) . Let σ be a norm continuous one-parameter group on A . Then φ is a KMS-weight with respect to σ if and only if*

1. *The mapping $\Lambda : \mathcal{N}_\varphi \mapsto H$ is closed.*
2. *We have that $\varphi \sigma_t = \varphi$ for every $t \in \mathbb{R}$.*
3. *There exists a core K for Λ such that*
 - *K is a subset of $D(\sigma_{\frac{i}{2}})$, $\sigma_{\frac{i}{2}}(K)^*$ is a subset of $\mathcal{N}_\varphi$ and $\|\Lambda(x)\| = \|\Lambda(\sigma_{\frac{i}{2}}(x)^*)\|$ for every $x \in K$.*
 - *We have that $\sigma_t(K) \subseteq K$ for every $t \in \mathbb{R}$.*

This proposition is a direct result of definition 5.11 and proposition 5.6. The next proposition is an easy consequence of definition 5.12 and proposition 5.14.

Proposition 6.2 *Consider a C^* -algebra A and a densely defined weight φ on A with GNS-construction (H, π, Λ) . Let σ be a norm continuous one-parameter group on A . Then φ is a KMS-weight with respect to σ if and only if*

1. *The mapping $\Lambda : \mathcal{N}_\varphi \mapsto H$ is closed.*

2. We have that $\varphi \sigma_t = \varphi$ for every $t \in \mathbb{R}$.

3. There exists an anti-unitary operator I on H such that the following holds.

There exists a core K for Λ and a core C for $\sigma_{\frac{i}{2}}$ such that we have for every $x \in K$ and $a \in C$ that xa belongs to $\mathcal{N}_\varphi$ and $\Lambda(xa) = I\pi(\sigma_{\frac{i}{2}}(a))^* I^* \Lambda(x)$.

Notice that the condition of lower semi-continuity of the weight is not assumed. This is replaced by the weaker condition of the closedness of the mapping Λ .

For the most part of this section, we will fix a C^* -algebra A and a KMS-weight φ on A with modular group σ . Let (H, Λ, π) be a GNS-construction for φ .

Because φ is invariant with respect to σ , there exists a unique injective positive operator ∇ on H such that $\nabla^{it} \Lambda(a) = \Lambda(\sigma_t(a))$ for every $a \in \mathcal{N}_\varphi$. Then we have that $\pi(\sigma_t(a)) = \nabla^{it} \pi(a) \nabla^{-it}$ for every $t \in \mathbb{R}$ and $a \in A$.

Define the set

$$C_\varphi = \{ a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \mid a \text{ is analytic with respect to } \sigma \text{ and } \sigma_z(a) \text{ belongs to } \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \text{ for every } z \in \mathbb{C} \} .$$

Then C_φ is a sub- $*$ -algebra of A . It is also clear that $\sigma_z(C) \subseteq C$ for every $z \in \mathbb{C}$.

Because φ is lower semi-continuous and densely defined, we know that $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ is a core for Λ . So, proposition 5.2 implies that C_φ is a core for $\mathcal{N}_\varphi$.

Let us also define the set

$$K_\varphi = \{ a \in \mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}}) \mid \sigma_{\frac{i}{2}}(a)^* \text{ belongs to } \mathcal{N}_\varphi \} .$$

Because, C_φ is a subset of K_φ , we have that K_φ is also a core for Λ . It is also clear that $\|\Lambda(a)\| = \|\Lambda(\sigma_{\frac{i}{2}}(a)^*)\|$ for every $a \in K_\varphi$. Furthermore, $\sigma_t(K_\varphi) \subseteq K_\varphi$ for every $t \in \mathbb{R}$.

Definition 6.3 Define the anti-unitary J on H such that $J\Lambda(a) = \Lambda(\sigma_{\frac{i}{2}}(a)^*)$ for every $a \in K_\varphi$.

We have that $JK_\varphi \subseteq K_\varphi$ and $J(J(v)) = v$ for every $v \in K_\varphi$. Therefore, $J^2 = 1$.

The proof of the following proposition is the same as the proof of proposition 5.9.

Proposition 6.4 Consider $x \in \mathcal{N}_\varphi$ and $a \in D(\sigma_{\frac{i}{2}})$. Then xa belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(xa) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x) .$$

In the same way as in the proof of lemma 5.10, the previous proposition and proposition 5.3 imply the following one.

Proposition 6.5 We have that φ is regular and has a truncating net consisting of elements of C_φ .

Result 6.6 Let k be a natural number. Then $(C_\varphi)^k$ is a core for Λ .

Proof : We know already that C_φ is a core for Λ . So we can suppose that $k \geq 2$. Choose $x \in C_\varphi$. Because $(C_\varphi)^{k-1}$ is dense in A , there exists a bounded net $(e_k)_{k \in K}$ in $(C_\varphi)^{k-1}$ such that $(e_k)_{k \in K}$ converges strictly to 1. Then we have for every $k \in K$ that $e_k x \in (C_\varphi)^k$. It is also clear that $(e_k x)_{k \in K}$ converges to x .
Because $\Lambda(e_k x) = \pi(e_k)\Lambda(x)$ for every $k \in K$, we get also that $(\Lambda(e_k x))_{k \in K}$ converges to $\Lambda(x)$.
Because C_φ is a core for Λ , this implies that $(C_\varphi)^k$ is a core for C_φ . ■

Combining this result with proposition 2.10, this implies the following result.

Result 6.7 *Let k be a natural number. Consider $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. Then there exists a sequence $(a_n)_{n=1}^\infty$ in $(C_\varphi)^k$ such that*

1. $(a_n)_{n=1}^\infty \rightarrow a$
2. $(\Lambda(a_n))_{n=1}^\infty \rightarrow \Lambda(a)$
3. $(\Lambda(a_n^*))_{n=1}^\infty \rightarrow \Lambda(a^*)$.

In the rest of this section, we will prove some familiar results

Proposition 6.8 *Consider $a \in D(\sigma_{-i})$ and $x \in \mathcal{M}_\varphi$. Then ax and $x\sigma_{-i}(a)$ belong to $\mathcal{M}_\varphi$ and $\varphi(ax) = \varphi(x\sigma_{-i}(a))$.*

Proof : Choose $y, z \in \mathcal{N}_\varphi$. Because a^* belongs to $D(\sigma_{\frac{i}{2}})$, we know that ya^* belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(ya^*) = J\pi(\sigma_{\frac{i}{2}}(a^*))^* J\Lambda(y) = J\pi(\sigma_{-\frac{i}{2}}(a))J\Lambda(y) .$$

Because $ay^*z = (ya^*)^*z$, we get that ay^*z belongs to $\mathcal{M}_\varphi$ and

$$\begin{aligned} \varphi(ay^*z) &= \varphi((ya^*)^*z) = \langle \Lambda(z), \Lambda(ya^*) \rangle \\ &= \langle \Lambda(z), J\pi(\sigma_{-\frac{i}{2}}(a))J\Lambda(y) \rangle = \langle J\pi(\sigma_{-\frac{i}{2}}(a))^* J\Lambda(z), \Lambda(y) \rangle \quad (*) \end{aligned}$$

We have that $\sigma_{-i}(a)$ belongs to $D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(\sigma_{-i}(a)) = \sigma_{-\frac{i}{2}}(a)$. This implies that $z\sigma_{-i}(a)$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(z\sigma_{-i}(a)) = J\pi(\sigma_{\frac{i}{2}}(\sigma_{-i}(a)))^* J\Lambda(z) = J\pi(\sigma_{-\frac{i}{2}}(a))^* J\Lambda(z) .$$

Therefore, $y^*z\sigma_{-i}(a)$ belongs to $\mathcal{M}_\varphi$ and

$$\varphi(y^*z\sigma_{-i}(a)) = \langle \Lambda(z\sigma_{-i}(a)), \Lambda(y) \rangle = \langle J\pi(\sigma_{-\frac{i}{2}}(a))^* J\Lambda(z), \Lambda(y) \rangle .$$

Comparing this with equation (*), we see that $\varphi(ay^*z) = \varphi(y^*z\sigma_{-i}(a))$. ■

Corollary 6.9 *Consider $a \in D(\sigma_i)$ and $x \in \mathcal{M}_\varphi$. Then xa and $\sigma_i(a)x$ belong to $\mathcal{M}_\varphi$ and $\varphi(xa) = \varphi(\sigma_i(a)x)$.*

Now, we prove a generalization of proposition 6.4.

Proposition 6.10 *Consider $x \in \mathcal{N}_\varphi$ and $a \in D(\overline{\sigma}_{\frac{i}{2}})$. Then xa belongs to $\mathcal{N}_\varphi$ and*

$$\Lambda(xa) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x) .$$

Proof : Choose $y \in \mathcal{N}_\varphi$ and $b \in D(\sigma_{\frac{i}{2}})$. By proposition 6.4, we know that yb belongs to $\mathcal{N}_\varphi$ and $\Lambda(yb) = J\pi(\sigma_{\frac{i}{2}}(b))^* J\Lambda(y)$.

We have also that ba belongs to $D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(ba) = \sigma_{\frac{i}{2}}(b)\sigma_{\frac{i}{2}}(a)$. Using proposition 6.4 once more, we see that $(yb)a$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda((yb)a) = J\pi(\sigma_{\frac{i}{2}}(ba))^* J\Lambda(y) = J\pi(\sigma_{\frac{i}{2}}(a))^* J J\pi(\sigma_{\frac{i}{2}}(b))^* J\Lambda(y) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(yb) .$$

Because $(C_\varphi)^2$ is a core for Λ , we have that $\langle yb \mid y \in \mathcal{N}_\varphi, b \in D(\sigma_{\frac{i}{2}}) \rangle$ is a core for Λ . Using the results of the previous discussion, this implies easily that xa belongs to $\mathcal{N}_\varphi$ and $\Lambda(xa) = J\pi(\sigma_{\frac{i}{2}}(a))^* J\Lambda(x)$. ■

Therefore, we can prove the next proposition in the same way as proposition 6.8.

Proposition 6.11 *Consider $a \in D(\overline{\sigma}_{-i})$ and $x \in \mathcal{M}_\varphi$. Then ax and $x\sigma_{-i}(a)$ belong to $\mathcal{M}_\varphi$ and $\varphi(ax) = \varphi(x\sigma_{-i}(a))$.*

Corollary 6.12 *Consider $a \in D(\overline{\sigma}_i)$ and $x \in \mathcal{M}_\varphi$. Then xa and $\sigma_i(a)x$ belong to $\mathcal{M}_\varphi$ and $\varphi(xa) = \varphi(\sigma_i(a)x)$.*

Another consequence of proposition 6.4 can be found in the following proposition.

Result 6.13 *Consider $a, b \in A$. Then we have that $J\pi(a)J$ and $\pi(b)$ commute.*

Proof : Choose $c \in D(\sigma_{\frac{i}{2}})$.

Take $x \in \mathcal{N}_\varphi$. Then xc belongs to $\mathcal{N}_\varphi$ and $\Lambda(xc) = J\pi(\sigma_{\frac{i}{2}}(c))^* J\Lambda(x)$. So bx belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(bxc) = \pi(b)\Lambda(xc) = \pi(b) J\pi(\sigma_{\frac{i}{2}}(c))^* J\Lambda(x) .$$

On the other hand, we have that bx belongs to $\mathcal{N}_\varphi$ and $\Lambda(bx) = \pi(b)\Lambda(x)$. This implies that bx belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(bxc) = J\pi(\sigma_{\frac{i}{2}}(c))^* J\Lambda(bx) = J\pi(\sigma_{\frac{i}{2}}(c))^* J\pi(b)\Lambda(x) .$$

Comparing these two expressions for $\Lambda(bxc)$, we see that

$$\pi(b) J\pi(\sigma_{\frac{i}{2}}(c))^* J\Lambda(x) = J\pi(\sigma_{\frac{i}{2}}(c))^* J\pi(b)\Lambda(x) .$$

This implies that $\pi(b) J\pi(\sigma_{\frac{i}{2}}(c))^* J = J\pi(\sigma_{\frac{i}{2}}(c))^* J\pi(b)$.

Because $\sigma_{\frac{i}{2}}$ has dense range in A , the result follows. ■

Proposition 6.14 *The set $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a core for $\nabla^{\frac{1}{2}}$ and $\Lambda(a^*) = J\nabla^{\frac{1}{2}}\Lambda(a)$ for every $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$.*

Proof : Choose $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. We know that there exists a sequence $a_n \in C_\varphi$ such that $(\Lambda(a_n))_{n=1}^\infty \rightarrow \Lambda(a)$ and $(\Lambda(a_n^*))_{n=1}^\infty \rightarrow \Lambda(a^*)$ (see result 6.7).

Choose $m \in \mathbb{N}$. Because a_m belongs to $\mathcal{N}_\varphi \cap D(\sigma_{-\frac{i}{2}})$ and $\sigma_{-\frac{i}{2}}(a_m)$ belongs to $\mathcal{N}_\varphi$, we have that $\Lambda(a_m)$ belongs to $D(\nabla^{\frac{1}{2}})$ and $\nabla^{\frac{1}{2}}\Lambda(a_m) = \Lambda(\sigma_{-\frac{i}{2}}(a_m))$.

We also know that a_m^* belongs to $\mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}})$ and $\sigma_{\frac{i}{2}}(a_m)^*$ belongs to $\mathcal{N}_\varphi^*$. By definition, we have that

$$J\Lambda(a_m^*) = \Lambda(\sigma_{\frac{i}{2}}(a_m^*)^*) = \Lambda(\sigma_{-\frac{i}{2}}(a_m)) = \nabla^{\frac{1}{2}}\Lambda(a_m) .$$

This implies that $(\nabla^{\frac{1}{2}}\Lambda(a_n))_{n=1}^\infty$ converges to $J\Lambda(a^*)$. The closedness of $\nabla^{\frac{1}{2}}$ implies that $\Lambda(a)$ belongs to $D(\nabla^{\frac{1}{2}})$ and $\nabla^{\frac{1}{2}}\Lambda(a) = J\Lambda(a^*)$.

Choose $x \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$, $n \in \mathbb{N}$ and define

$$x(n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \sigma_t(x) dt ,$$

it is clear that $x(n)$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ and

$$\Lambda(x(n)) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \nabla^{it} \Lambda(x) dt .$$

Because $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is dense in H , this implies that $\langle \Lambda(x(n)) \mid x \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*, n \in \mathbb{N} \rangle$ is a core for $\nabla^{\frac{1}{2}}$. This implies that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a core for $\nabla^{\frac{1}{2}}$. $\blacksquare$

We want to construct a left Hilbert algebra in the usual way. First, we will need a lemma which guarantees that our next definitions are well defined.

Lemma 6.15 *1. Consider $a_1, a_2, b_1, b_2 \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ such that $\Lambda(a_1) = \Lambda(a_2)$ and $\Lambda(b_1) = \Lambda(b_2)$. Then $\Lambda(a_1 b_1) = \Lambda(a_2 b_2)$.*

2. Consider $a_1, a_2 \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ such that $\Lambda(a_1) = \Lambda(a_2)$. Then $\Lambda(a_1^*) = \Lambda(a_2^*)$.

Proof :

1. For any $c \in C_\varphi$, we have that

$$\pi(a_1) \Lambda(c) = J \pi(\sigma_{\frac{i}{2}}(c))^* J \Lambda(a_1) = J \pi(\sigma_{\frac{i}{2}}(c))^* J \Lambda(a_2) = \pi(a_2) \Lambda(c) .$$

The density of $\Lambda(C_\varphi)$ in H implies that $\pi(a_1) = \pi(a_2)$.

Therefore,

$$\Lambda(a_1 b_1) = \pi(a_1) \Lambda(b_1) = \pi(a_2) \Lambda(b_2) = \Lambda(a_2 b_2) .$$

2. This follows easily from the previous proposition. $\blacksquare$

Definition 6.16 *We define $\mathcal{U} = \Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$, then $\mathcal{U}$ is subspace of H . We make $\mathcal{U}$ into a $*$ -algebra such that :*

1. We have for every $a, b \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ that $\Lambda(ab) = \Lambda(a) \Lambda(b)$.

2. We have for every $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ that $\Lambda(a)^* = \Lambda(a^*)$.

Then $\mathcal{U}$ becomes a left Hilbert algebra on H .

It is not difficult to check that $\mathcal{U}$ satisfies the conditions of a left Hilbert algebra. The closability of the mapping $\mathcal{U} \rightarrow \mathcal{U} : v \mapsto v^*$ is a direct consequence of proposition 6.14.

It is immediately clear that $L_{\Lambda(a)} = \pi(a)$ for every $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$.

Let us define T as the closure of the mapping $\mathcal{U} \rightarrow \mathcal{U} : v \mapsto v^*$. Proposition 6.14 implies that $T = J \nabla^{\frac{1}{2}}$.

So we see that J is the modular conjugation of $\mathcal{U}$ and ∇ is the modular operator of $\mathcal{U}$.

We also see that J and ∇ are independent of our choice of σ . Therefore, we will call ∇ the modular operator of φ and J the modular conjugation of φ in the GNS-construction (H, Λ, π) .

Lemma 6.17 *Consider $a \in \mathcal{N}_\varphi$. Then $\Lambda(a)$ is left bounded with respect to $\mathcal{U}$ and $L_{\Lambda(a)} = \pi(a)$.*

Proof : Because $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ is a core for Λ , there exists a sequence $(a_n)_{n=1}^\infty$ in $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ such that $(a_n)_{n=1}^\infty$ converges to a and $(\Lambda(a_n))_{n=1}^\infty$ converges to $\Lambda(a)$.

We know already that $L_{\Lambda(a_n)} = \pi(a_n)$ for $n \in \mathbb{N}$. This implies that $(L_{\Lambda(a_n)})_{n=1}^\infty$ converges to $\pi(a)$. Consequently, we find that $\Lambda(a)$ is left bounded with respect to $\mathcal{U}$ and $L_{\Lambda(a)} = \pi(a)$. ■

Lemma 6.18 *Consider $v \in H$ such that v is left bounded with respect to $\mathcal{U}$ and such that there exists an element $a \in A$ such that $L_v = \pi(a)$. Then a belongs to $\mathcal{N}_\varphi$ and $\Lambda(a) = v$.*

Proof : Choose a truncating net $(u_i)_{i \in I}$ for φ . For every $i \in I$, we define the operator $S_i \in B(H)$ such that $S_i \Lambda(x) = \Lambda(xu_i)$ for every $x \in A$.

By [6], there exist a sequence $(a_n)_{n=1}^\infty$ in $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ such that $(\Lambda(a_n))_{n=1}^\infty$ converges to $\Lambda(a)$, $(L_{\Lambda(a_n)})_{n=1}^\infty$ converges strongly to L_v and $(L_{\Lambda(a_n)})_{n=1}^\infty$ is bounded.

Choose $j \in I$. We have immediately that $(L_{\Lambda(a_n)} \Lambda(u_j))_{n=1}^\infty$ converges to $L_v \Lambda(u_j)$. We have for every $n \in \mathbb{N}$ that

$$L_{\Lambda(a_n)} \Lambda(u_j) = \pi(a_n) \Lambda(u_j) = \Lambda(a_n u_j) = S_j \Lambda(a_n) .$$

This implies that $(L_{\Lambda(a_n)} \Lambda(u_j))_{n=1}^\infty$ converges to $S_j v$.

Combining these two results, we get that $L_v \Lambda(u_j) = S_j v$, hence

$$\Lambda(a u_j) = \pi(a) \Lambda(u_j) = L_v \Lambda(u_j) = S_j v .$$

Therefore, $(\Lambda(a u_i))_{i \in I}$ converges to v . We also have that $(a u_i)_{i \in I}$ converges to a . The closedness of Λ implies that a belongs to $\mathcal{N}_\varphi$ and $\Lambda(a) = v$. ■

Lemma 6.19 *Consider $v \in \mathcal{U}''$ such that there exist an element $a \in A$ such that $L_v = \pi(a)$. Then a belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ and $v = \Lambda(a)$.*

Proof : We know that v is left bounded. Therefore, the previous lemma implies that a belongs to $\mathcal{N}_\varphi$ and $v = \Lambda(a)$. We have also that v^* is left bounded and $L_{v^*} = (L_v)^* = \pi(a)^* = \pi(a^*)$. Hence, the previous lemma implies that a^* belongs to $\mathcal{N}_\varphi$. So we get that a belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. ■

Using this result, we have the following theorem. This is a well known result for weights satisfying the other KMS-condition (proved by Combes, see lemma 2.2.3 of [5]).

Theorem 6.20 *Call $\tilde{\varphi}$ the normal semi-finite faithful weight on $\pi(A)''$ associated to $\mathcal{U}$. Then $\tilde{\varphi} \pi = \varphi$.*

Remark 6.21 Consider the case where we get a KMS-weight from a left Hilbert algebra (the third part of section 5, page 34) :

Consider a Hilbert space H and a left Hilbert $\mathcal{W}$ algebra on H with modular operator Δ , modular conjugation I and associated faithful semi-finite normal weight ψ .

Let A be a C^* -algebra and π be a nondegenerate $*$ -representation of A on H such that $\pi(A) \subseteq \mathcal{L}(\mathcal{W})$ and suppose there exists a one-parameter group σ on A such that $\pi(\sigma_t(a)) = \Delta^{it} \pi(a) \Delta^{-it}$ for every $t \in \mathbb{R}$ and $a \in A$.

Put $\varphi = \psi \pi$, so φ is a lower semi-continuous weight on A .

Define the following mapping Λ from $\mathcal{N}_\varphi$ into H : For every $a \in \mathcal{N}_\varphi$, there exists a unique element $v \in H$ such that v is left bounded with respect to $\mathcal{W}$ and $L_v = \pi(a)$ and we define $\Lambda(a) = v$, so we get that $L_{\Lambda(a)} = \pi(a)$. We suppose that $\mathcal{N}_\varphi$ is dense in A and that $\Lambda(\mathcal{N}_\varphi)$ is dense in H .

We saw that φ is a KMS-weight on A with modular group σ and GNS-construction (H, Λ, π) . Now we can apply the construction procedures of this section to get the left Hilbert algebra $\mathcal{U}$, the weight $\tilde{\varphi}$ associated to $\mathcal{U}$, the modular operator ∇ and modular conjugation J of φ in the GNS-construction (H, Λ, π) .

Proposition 5.16 implies that $\mathcal{U}'' = \mathcal{W}''$. Therefore we get that $\tilde{\varphi} = \psi$, $\nabla = \Delta$ and $J = I$.

For KMS-weights, the faithfulness of φ and π are equivalent :

Proposition 6.22 *We have that φ is faithful if and only if π is faithful.*

Proof :

$\Rightarrow$ This follows easily because φ is densely defined.

$\Leftarrow$ Choose $x \in A$ such that $\varphi(x^*x) = 0$. Then x belongs to $\mathcal{N}_\varphi$ and $\Lambda(x) = 0$.

Take $m \in \mathbb{N}$ and define

$$x(m) = \frac{m}{\sqrt{\pi}} \int \exp(-m^2 t^2) \sigma_t(x) dt .$$

We have that $x(m)$ belongs to $\mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}})$ and

$$\Lambda(x(m)) = \frac{m}{\sqrt{\pi}} \int \exp(-m^2 t^2) \nabla^{it} \Lambda(x) dt = 0 .$$

So we have for every $a \in \mathcal{N}_\varphi$ that

$$J\pi(\sigma_{\frac{i}{2}}(x(m)))^* J\Lambda(a) = \pi(a)\Lambda(x(m)) = 0 ,$$

which implies that $\pi(\sigma_{\frac{i}{2}}(x(m))) = 0$. The faithfulness of π implies that $\sigma_{\frac{i}{2}}(x(m)) = 0$, the injectivity of $\sigma_{\frac{i}{2}}$ implies that $x(m) = 0$.

The definition of the elements $x(n)$ ($n \in \mathbb{N}$) implies that $(x(n))_{n=1}^\infty \rightarrow x$. Therefore we have that $x = 0$. ■

We have of course also the following result.

Lemma 6.23 *Consider a $*$ -automorphism α on A such that there exists a strictly positive number λ such that $\varphi \alpha = \lambda \varphi$. Define u as the unitary operator on H such that $u\Lambda(a) = \lambda^{-\frac{1}{2}} \Lambda(\alpha(a))$ for every $a \in \mathcal{N}_\varphi$. Then $u\nabla = \nabla u$ and $uJ = Ju$. We have moreover that $\pi(\alpha(a)) = u\pi(a)u^*$ for every $a \in A$.*

Proof : Because α is a $*$ -homomorphism, it is easy to check for every $a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ that $u\Lambda(a)$ belongs to $D(T)$ and $Tu\Lambda(a) = uT\Lambda(a)$. Because $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ is a core for T and $u\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*) = \Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*)$ we get that $uT = Tu$. So we get also that $Tu^* = u^*T$. The unitarity of u implies that

$$uT^* = (Tu^*)^* = (u^*T)^* = T^*u .$$

Using the fact that $\nabla = T^*T$, this implies that $u\nabla = \nabla u$.

So we get also that $u\nabla^{\frac{1}{2}} = \nabla^{\frac{1}{2}}u$. This in turn implies that

$$uJ\nabla^{\frac{1}{2}} = uT = Tu = J\nabla^{\frac{1}{2}}u = Ju\nabla^{\frac{1}{2}} .$$

Because $\nabla^{\frac{1}{2}}$ has dense range, this implies immediately that $uJ = Ju$. ■

Corollary 6.24 *Suppose that φ is faithful. Consider a $*$ -automorphism α on A such that there exists a strictly positive number λ such that $\varphi \alpha = \lambda \varphi$. Then $\alpha \sigma_t = \sigma_t \alpha$ for every $t \in \mathbb{R}$.*

Proof : Choose $t \in \mathbb{R}$.

Define u as the unitary operator on H such that $u\Lambda(a) = \lambda^{-\frac{1}{2}} \Lambda(\alpha(a))$ for every $a \in \mathcal{N}_\varphi$. Then the previous lemma implies that $u\nabla = \nabla u$.

This implies immediately that $u\nabla^{it} = \nabla^{it}u$. So we get for every $a \in \mathcal{N}_\varphi$ that

$$\Lambda(\alpha(\sigma_t(a))) = \lambda^{\frac{1}{2}} u\Lambda(\sigma_t(a)) = \lambda^{\frac{1}{2}} u\nabla^{it}\Lambda(a) = \lambda^{\frac{1}{2}} \nabla^{it}u\Lambda(a) = \nabla^{it}\Lambda(\alpha(a)) = \Lambda(\sigma_t(\alpha(a))) .$$

so the faithfulness of φ implies that $\alpha(\sigma_t(a)) = \sigma_t(\alpha(a))$. ■

From the theory of left Hilbert algebras (see e.g. [18]), we get the following result :

Proposition 6.25 *The weight φ satisfies the KMS-condition with respect to σ : For every $x, y \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ there exists a bounded continuous function f from the strip $S(i)$ into $\mathbb{C}$ which is analytic on $S(i)^0$ and satisfies :*

- $f(t) = \varphi(\sigma_t(x)y)$ for all $t \in \mathbb{R}$
- $f(t+i) = \varphi(x\sigma_t(y))$ for all $t \in \mathbb{R}$.

The next two results follow immediately from proposition 6.10 and its corollary.

Proposition 6.26 *Consider $a \in M(A)$ such that there exists a strictly positive number λ such that $\sigma_t(a) = \lambda^{it} a$ for every $t \in \mathbb{R}$. Let x be an element in $\mathcal{N}_\varphi$. Then xa belongs to $\mathcal{N}_\varphi$ and $\Lambda(xa) = \lambda^{-\frac{1}{2}} J\pi(a)^* J\Lambda(x)$.*

Proposition 6.27 *Consider $a \in M(A)$ such that there exists a strictly positive number λ such that $\sigma_t(a) = \lambda^{it} a$ for every $t \in \mathbb{R}$. Let x be an element in $\mathcal{M}_\varphi$. Then ax and xa belong to $\mathcal{M}_\varphi$ and $\varphi(ax) = \lambda\varphi(xa)$.*

If φ is faithful, it is possible to prove a converse of this corollary.

The proof of the next lemma can be copied immediately from the proof of lemma 3.4 of [14].

Lemma 6.28 *Consider $a \in M(A)$ such that $a\mathcal{M}_\varphi \subseteq \mathcal{M}_\varphi$ and $\mathcal{M}_\varphi a \subseteq \mathcal{M}_\varphi$. Then $\sigma_t(a)\mathcal{M}_\varphi \subseteq \mathcal{M}_\varphi$ and $\mathcal{M}_\varphi \sigma_t(a) \subseteq \mathcal{M}_\varphi$ for every $t \in \mathbb{R}$. Let x, y be elements in C_φ , then there exists an entire function from $\mathbb{C}$ into $\mathbb{C}$ such that*

1. *We have that $f(t) = \varphi(\sigma_t(a)y^*x)$ for every $t \in \mathbb{R}$.*
2. *We have that $f(t+i) = \varphi(y^*x\sigma_t(a))$ for every $t \in \mathbb{R}$.*
3. *f is bounded on each horizontal strip.*

The proof of the following proposition is a slight adaptation of the proof of theorem 3.6 of [14].

Result 6.29 *Suppose that φ is faithful. Consider $a \in M(A)$ such that $a\mathcal{M}_\varphi \subseteq \mathcal{M}_\varphi$, $\mathcal{M}_\varphi a \subseteq \mathcal{M}_\varphi$ and such that there exist a strictly positive number λ satisfying $\varphi(ax) = \lambda\varphi(xa)$ for every $x \in \mathcal{M}_\varphi$. Then $\sigma_t(a) = \lambda^{it} a$ for every $t \in \mathbb{R}$.*

Proof : Choose $x, y \in C_\varphi$.

By the previous lemma, there exists an entire function f from $\mathbb{C}$ into $\mathbb{C}$ such that

1. We have that $f(t) = \varphi(\sigma_t(a)y^*x)$ for every $t \in \mathbb{R}$.

2. We have that $f(t+i) = \varphi(y^*x\sigma_t(a))$ for every $t \in \mathbb{R}$.
3. f is bounded on each horizontal strip.

First, we prove that $f(z+i) = \lambda^{-1}f(z)$ for every $z \in \mathbb{C}$.

We have for every $t \in \mathbb{R}$ that

$$\begin{aligned}
f(t+i) - \lambda^{-1}f(t) &= \varphi(y^*x\sigma_t(a)) - \lambda^{-1}\varphi(\sigma_t(a)y^*x) \\
&= \varphi(y^*x\sigma_t(a)) - \lambda^{-1}\varphi(a\sigma_{-t}(y^*x)) \\
&= \varphi(y^*x\sigma_t(a)) - \lambda^{-1}\lambda\varphi(\sigma_{-t}(y^*x)a) \\
&= \varphi(y^*x\sigma_t(a)) - \varphi(y^*x\sigma_t(a)) = 0.
\end{aligned}$$

By analyticity of the function

$$\mathbb{C} \rightarrow \mathbb{C} : z \mapsto f(z+i) - \lambda^{-1}f(z),$$

we must have that $f(z+i) - \lambda^{-1}f(z) = 0$ for every $z \in \mathbb{C}$.

Now we define the entire function g from $\mathbb{C}$ into $\mathbb{C}$ such that $g(z) = \lambda^{-iz}f(z)$ for every $z \in \mathbb{C}$. Then we have for every $z \in \mathbb{C}$ that

$$g(z+i) = \lambda^{-i(z+i)}f(z+i) = \lambda^{-iz}\lambda\lambda^{-1}f(z) = \lambda^{-iz}f(z) = g(z). \quad (*)$$

We know that f is bounded on $S(i)$. We have also that

$$|g(z)| = |\lambda^{-iz}| |f(z)| = \lambda^{\text{Im } z} |f(z)| \leq \max\{1, \lambda\} |f(z)|,$$

for every $z \in S(i)$, so g is also bounded on $S(i)$.

By (*), we have that g is bounded on $\mathbb{C}$. The theorem of Liouville implies that g is a constant function. This implies that

$$\varphi(ay^*x) = f(0) = g(0) = g(t) = \lambda^{-it}f(t) = \lambda^{-it}\varphi(\sigma_t(a)y^*x)$$

for every $t \in \mathbb{R}$.

Choose $s \in \mathbb{R}$. Because $(a - \lambda^{-is}\sigma_s(a))\mathcal{M}_\varphi \subseteq \mathcal{M}_\varphi$, we have also that $\mathcal{N}_\varphi(a^* - \lambda^{is}\sigma_s(a^*)) \subseteq \mathcal{N}_\varphi$. Hence, we have for every $x, y \in C_\varphi$ that

$$\langle \Lambda(x), \Lambda(y(a^* - \lambda^{is}\sigma_s(a^*))) \rangle = \varphi((a - \lambda^{-is}\sigma_s(a))y^*x) = 0.$$

Because $\Lambda(C_\varphi)$ is dense in H , we see that $\Lambda(y(a^* - \lambda^{is}\sigma_s(a^*))) = 0$ for every $y \in C_\varphi$. The faithfulness of φ implies that $y(a^* - \lambda^{is}\sigma_s(a^*)) = 0$ for every $y \in C_\varphi$. Therefore, the density of C_φ in A implies that $a^* - \lambda^{is}\sigma_s(a^*) = 0$. ■

The following proposition is an immediate consequence of proposition 6.26.

Proposition 6.30 *Consider a strictly positive element $\delta \in A$ such that there exist a strictly positive number λ such that $\sigma_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$. Then we have the following properties :*

- For every $x \in \mathcal{N}_\varphi$ and every $t \in \mathbb{R}$, we have that $x\delta^{it}$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(x\delta^{it}) = \lambda^{-\frac{1}{2}} J\pi(\delta)^{-it} J\Lambda(x)$.
- For every $x \in \mathcal{M}_\varphi$ and every $t \in \mathbb{R}$, we have that $x\delta^{it}$ and $\delta^{it}x$ belong to $\mathcal{M}_\varphi$ and $\varphi(\delta^{it}x) = \lambda^t \varphi(x\delta^{it})$.

Using this proposition, we get the following one.

Proposition 6.31 *Consider a strictly positive element $\delta \in A$ such that there exist a strictly positive number λ such that $\sigma_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$. Let z be a complex number and $a \in \mathcal{N}_\varphi$ such that a is a left multiplier of δ^z and $a\delta^z$ belongs to A . Then we have the following properties :*

- $a\delta^z$ belongs to $\mathcal{N}_\varphi \Leftrightarrow \Lambda(a)$ belongs to $D(J\pi(\delta)^{\bar{z}}J)$.
- If $a\delta^z$ belongs to $\mathcal{N}_\varphi$, then $\Lambda(a\delta^z) = \lambda^{\frac{iz}{2}} J\pi(\delta)^{\bar{z}}J\Lambda(a)$.

Proof : We define the strongly continuous one-parameter representation α on A such that $\alpha_t(x) = x\delta^{it}$ for every $x \in A$ and $t \in \mathbb{R}$.

Furthermore, we define the strongly continuous unitary group representation u from $\mathbb{R}$ on H such that $u_t = (J\pi(\delta)J)^{it}$ for every $t \in \mathbb{R}$. We know by the previous proposition that

1. We have that $\alpha_t(\mathcal{N}_\varphi) \subseteq \mathcal{N}_\varphi$ for every $t \in \mathbb{R}$
2. We have that $u_t\Lambda(a) = \lambda^{\frac{t}{2}} \Lambda(\alpha_t(a))$ for every $t \in \mathbb{R}$ and $a \in \mathcal{N}_\varphi$.

Because a belongs to $D(\alpha_{-iz})$ and $\alpha_{-iz}(a) = a\delta^z$ and $u_{-iz} = (J\pi(\delta)J)^z$, this proposition follows from propositions 4.4 and 4.5. ■

Proposition 6.32 *Consider a strictly positive element δ affiliated with A such that there exists a strictly positive number λ such that $\sigma_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$. Let a be an element in A and $z \in \mathbb{C}$ such that a is a left and right multiplier of δ^z and $a\delta^z, \delta^z a$ belong to $\mathcal{M}_\varphi$. Then $\varphi(a\delta^z) = \lambda^{iz} \varphi(\delta^z a)$.*

Proof : Define the norm continuous one-parameter group α on A such that $\alpha_t(a) = \delta^{it} a \delta^{-it}$ for every $a \in A$ and $t \in \mathbb{R}$. Proposition 6.30 implies that $\varphi \alpha_t = \lambda^t \varphi$ for every $t \in \mathbb{R}$.

Because a is a left multiplier of δ^z , we have that $a\delta^z$ is a left multiplier of δ^{-z} and $(a\delta^z)\delta^{-z} = a$.

This implies that $(a\delta^z)\delta^{-z}$ is a right multiplier of δ^z and $\delta^z [(a\delta^z)\delta^{-z}] = \delta^z a$. From this, we get that $a\delta^z$ is a middle multiplier of δ^z, δ^{-z} and

$$\delta^z (a\delta^z) \delta^{-z} = \delta^z a$$

From this all, we get that $a\delta^z$ belongs to $D(\alpha_{-iz})$ and $\alpha_{-iz}(a\delta^z) = \delta^z a$. The proposition follows from proposition 2.14. ■

We now prove that a KMS-weight has no proper extensions which are invariant under its modular group.

Proposition 6.33 *Consider a lower semi-continuous weight η on A which is an extension of φ and such that η is invariant under σ . Then $\varphi = \eta$.*

Proof : Take a GNS-construction $(H_\eta, \Lambda_\eta, \pi_\eta)$ for η .

Because η is invariant under σ , we get the existence a positive injective operator T in H_η such that $T^{it}\Lambda_\eta(a) = \Lambda_\eta(\sigma_t(a))$ for every $a \in \mathcal{N}_\eta$ and $t \in \mathbb{R}$.

Choose $y \in \mathcal{N}_\eta$.

Define for every $n \in \mathbb{N}$ the element

$$y_n = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \sigma_t(y) dt$$

which is clearly analytic with respect to σ . We have also that $(y_n)_{n=1}^\infty$ converges to y .

By ??, we have for every $n \in \mathbb{N}$ that y_n belongs to $\mathcal{N}_\eta$ and

$$\Lambda_\eta(y_n) = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) T^{it} \Lambda(y) dt .$$

This implies immediately that $(\Lambda_\eta(y_n))_{n=1}^\infty$ converges to $\Lambda_\eta(y)$.

We can also take an approximate unit $(e_i)_{i \in I}$ for A in $\mathcal{N}_\varphi$. Then $(e_i y_n)_{(i,n) \in I \times \mathbb{N}}$ converges to y .

We have also for every $i \in I$ and $n \in \mathbb{N}$ that $e_i y_n$ belongs to $\mathcal{N}_\eta$ and $\Lambda_\eta(e_i y_n) = \pi_\eta(e_i) \Lambda_\eta(y_n)$.

Consequently, the net $(\Lambda_\eta(e_i y_n))_{(i,n) \in I \times \mathbb{N}}$ converges to $\Lambda_\eta(y)$.

We have for every $i \in I$ and $n \in \mathbb{N}$ that e_i belongs to $\mathcal{N}_\varphi$ and y_n belongs to $D(\sigma_{\frac{i}{2}})$ implying that $e_i y_n$ belongs to $\mathcal{N}_\varphi$ by proposition 6.4.

Because $\varphi \subseteq \eta$, we have moreover for every $i, j \in I$ and $m, n \in \mathbb{N}$ that

$$\|\Lambda_\varphi(e_i y_n) - \Lambda_\varphi(e_j y_m)\| = \|\Lambda_\eta(e_i y_n) - \Lambda_\eta(e_j y_m)\| .$$

This last equality implies that the net $(\Lambda_\varphi(e_i y_n))_{(i,n) \in I \times \mathbb{N}}$ is Cauchy and hence convergent in H_φ . Therefore, the closedness of Λ_φ implies that y is an element of $\mathcal{N}_\varphi$. The proposition follows. $\blacksquare$

The following proposition will guarantee that the modular group is unique for faithful KMS-weights.

Proposition 6.34 *Consider a C^* -algebra A and a KMS-weight φ on A with GNS-construction (H, Λ, π) . Let σ and τ be modular groups for φ . Then $\pi \sigma_t = \pi \tau_t$ for every $t \in \mathbb{R}$.*

Proof : Define the injective positive operators ∇ and Δ in H such that $\nabla^{it} \Lambda(a) = \Lambda(\sigma_t(a))$ and $\Delta^{it} \Lambda(a) = \Lambda(\tau_t(a))$ for every $a \in \mathcal{N}_\varphi$ and $t \in \mathbb{R}$. Define also the anti-unitary operators J, I on H such that

- $J\Lambda(a) = \Lambda(\sigma_{\frac{i}{2}}(a)^*)$ for every $a \in \mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}})$ such that $\sigma_{\frac{i}{2}}(a)^*$ belongs to $\mathcal{N}_\varphi$
- $I\Lambda(a) = \Lambda(\tau_{\frac{i}{2}}(a)^*)$ for every $a \in \mathcal{N}_\varphi \cap D(\tau_{\frac{i}{2}})$ such that $\tau_{\frac{i}{2}}(a)^*$ belongs to $\mathcal{N}_\varphi$

Proposition 6.14 implies that $J\nabla^{\frac{1}{2}} = I\Delta^{\frac{1}{2}}$. The uniqueness of the polar decomposition implies that $\nabla = \Delta$. This implies for every $t \in \mathbb{R}$ that

$$\pi(\sigma_t(a)) = \nabla^{it} \pi(a) \nabla^{-it} = \Delta^{it} \pi(a) \Delta^{-it} = \pi(\tau_t(a))$$

for every $t \in \mathbb{R}$ and $a \in A$. $\blacksquare$

Corollary 6.35 *Consider a C^* -algebra A and a faithful KMS-weight φ on A . Then φ as a unique modular group.*

In the last theorem of this section, we prove that our definition of KMS-weight is equivalent with the usual one (introduced in [4]).

Theorem 6.36 *Consider a C^* -algebra A , a densely defined lower semi-continuous weight φ on A and a one-parameter group σ on A . Then φ is a KMS-weight with respect to σ if and only if*

1. We have for every $t \in \mathbb{R}$ that $\varphi \sigma_t = \varphi$.
2. For every $x, y \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$, there exists a bounded continuous function f from $S(i)$ into $\mathbb{C}$ which is analytic on $S(i)^0$ and such that :
 - We have for every $t \in \mathbb{R}$ that $f(t) = \varphi(\sigma_t(x)y)$.
 - We have for every $t \in \mathbb{R}$ that $f(t+i) = \varphi(y\sigma_t(x))$.

Proof :

$\Rightarrow$ This follows from proposition 6.25.

$\Leftarrow$ Take a GNS-construction (H, Λ, π) for φ . Define the injective positive operator ∇ in H such that $\nabla^{it} \Lambda(a) = \Lambda(\sigma_t(a))$ for every $t \in \mathbb{R}$.

Define the set

$$C = \{ a \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \mid a \text{ is analytic with respect to } \sigma \text{ and } \sigma_z(a) \text{ belongs to } \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \text{ for every } z \in \mathbb{C} \} .$$

So C is a sub*-algebra of A such that $\sigma_z(C) \subseteq C$ for every $z \in \mathbb{C}$.

Choose $x, y \in C$. Because x, y belong to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$, we have by assumption the existence of a bounded continuous function f on $S(i)$ which is analytic on $S(i)^0$ and satisfies :

- We have for every $t \in \mathbb{R}$ that $f(t) = \varphi(\sigma_t(x)y)$.
- We have for every $t \in \mathbb{R}$ that $f(t+i) = \varphi(y\sigma_t(x))$.

We know that x^* belongs to $\mathcal{N}_\varphi$ and that $\sigma_{-i}(x^*)$ belongs to $\mathcal{N}_\varphi$. By proposition 4.4, we get that $\Lambda(x^*)$ belongs to $D(\nabla)$ and $\nabla \Lambda(x^*) = \Lambda(\sigma_{-i}(x^*))$. Define the function g from $S(i)$ into $\mathbb{C}$ such that $g(z) = \langle \Lambda(y), \nabla^{i\bar{z}} \Lambda(x^*) \rangle$ for every $z \in S(i)$. Then g is continuous on $S(i)$ and analytic on $S(i)^0$. We have for every $t \in \mathbb{R}$ that

$$g(t) = \langle \Lambda(y), \nabla^{it} \Lambda(x^*) \rangle = \langle \Lambda(y), \Lambda(\sigma_t(x^*)) \rangle = \varphi(\sigma_t(x)y) = f(t) .$$

Therefore we must have that $f = g$ which implies that

$$\varphi(yx) = f(i) = g(i) = \langle \Lambda(y), \nabla \Lambda(x^*) \rangle = \langle \Lambda(y), \Lambda(\sigma_{-i}(x^*)) \rangle = \varphi(\sigma_i(x)y) . \quad (*)$$

We know already that $C \subseteq \mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}})$ and that $\sigma_{\frac{i}{2}}(C)^* \subseteq \mathcal{N}_\varphi$. Just like in lemma 5.13, we get that C is a core for Λ . Choose $a \in C$. If we replace in equation (*) the element x by $\sigma_{-\frac{i}{2}}(a)$ and y by $\sigma_{\frac{i}{2}}(a)^*$, we get that

$$\varphi(\sigma_{\frac{i}{2}}(a)^* \sigma_{-\frac{i}{2}}(a)) = \varphi(\sigma_{\frac{i}{2}}(a) \sigma_{\frac{i}{2}}(a)^*) .$$

We have that $\sigma_{-\frac{i}{2}}(a^*a) = \sigma_{\frac{i}{2}}(a)^* \sigma_{-\frac{i}{2}}(a)$ which belongs to $\mathcal{M}_\varphi$. On the other hand, we know also that a^*a belongs to $\mathcal{M}_\varphi$. Hence, proposition 2.14 implies that

$$\varphi(\sigma_{\frac{i}{2}}(a)^* \sigma_{-\frac{i}{2}}(a)) = \varphi(\sigma_{\frac{i}{2}}(a^*a)) = \varphi(a^*a) .$$

Combining these two equalities, we get that $\varphi(a^*a) = \varphi(\sigma_{\frac{i}{2}}(a) \sigma_{\frac{i}{2}}(a)^*)$. Therefore, proposition 6.1 implies that φ is KMS with respect to σ .

■

In order to prove the implication from the right to the left, we do not really need the lower semi-continuity of φ . The closedness of the map $\Lambda : \mathcal{N}_\varphi \rightarrow H$ is also sufficient.

7 The tensor product of KMS-weights

Consider two C^* -algebras A and B . Let φ be a KMS-weight on A with modular group σ and ψ a KMS-weight on B with modular group τ .

Take also a GNS-construction (H, Λ, π) for φ and a GNS-construction (K, Γ, θ) for ψ .

- With respect to the GNS-construction (H, Λ, π) , we denote the modular conjugation of φ by J and the modular operator of φ by ∇ .
- With respect to the GNS-construction (K, Γ, θ) , we denote the modular conjugation of ψ by I and the modular operator ψ by Δ .

We will define the weight $\varphi \otimes \psi$ and show that it is a KMS-weight with respect to the obvious one-parameter group. The construction of the weight itself is due to Jan Verding and can be found in [22]. He did it in fact in a more general case. Again, due to the lack of availability of this work, we will include the proofs. We only added the KMS-characteristics.

Definition 7.1 *We define the norm continuous one-parameter group $\sigma \otimes \tau$ on $A \otimes B$ such that $(\sigma \otimes \tau)_t = \sigma_t \otimes \tau_t$ for every $t \in \mathbb{R}$.*

Consider $z \in \mathbb{C}$. Then it is easy to see that $\sigma_z \odot \tau_z \subseteq (\sigma \otimes \tau)_z$. We will prove in [9] (and this is not too difficult, using smearing techniques) that $D(\sigma_z) \odot D(\tau_z)$ is a core for $(\sigma \otimes \tau)_z$. But we will not need this fact in this paper.

Lemma 7.2 *The mapping $\Lambda \odot \Gamma : \mathcal{N}_\varphi \odot \mathcal{N}_\psi \mapsto H \otimes K$ is closable.*

Proof : Define for every $\omega \in \mathcal{F}_\varphi$ the operator $S_\omega \in B(H)$ such that $\langle S_\omega \Lambda(a), \Lambda(b) \rangle = \omega(b^*a)$ for every $a, b \in \mathcal{N}_\varphi$. In a similar way, we define for every $\eta \in \mathcal{F}_\psi$ the operator $T_\eta \in B(K)$ such that $\langle T_\eta \Gamma(a), \Gamma(b) \rangle = \eta(b^*a)$ for every $a, b \in \mathcal{N}_\psi$.

Take a sequence $(x_n)_{n=1}^\infty$ in $\mathcal{N}_\varphi \odot \mathcal{N}_\psi$ and $v \in H \otimes K$ such that $(x_n)_{n=1}^\infty$ converges to 0 and $((\Lambda \odot \Gamma)(x_n))_{n=1}^\infty$ converges to v .

Fix $\omega \in \mathcal{F}_\varphi$ and $\eta \in \mathcal{F}_\psi$ for the moment.

Choose $a \in \mathcal{N}_\varphi$ and $b \in \mathcal{N}_\psi$. It is not difficult to see that

$$\langle (S_\omega \otimes T_\eta)(\Lambda \otimes \Gamma)(x_n), \Lambda(a) \otimes \Gamma(b) \rangle = (\omega \otimes \eta)((a \otimes b)^* x_n)$$

for every $n \in \mathbb{N}$. This implies that the sequence

$$\left(\langle (S_\omega \otimes T_\eta)(\Lambda \otimes \Gamma)(x_n), \Lambda(a) \otimes \Gamma(b) \rangle \right)_{n=1}^\infty$$

converges to 0. It is also clear that the sequence

$$\left(\langle (S_\omega \otimes T_\eta)(\Lambda \otimes \Gamma)(x_n), \Lambda(a) \otimes \Gamma(b) \rangle \right)_{n=1}^\infty$$

converges to $\langle (S_\omega \otimes T_\eta)v, \Lambda(a) \otimes \Gamma(b) \rangle$. So we see that $\langle (S_\omega \otimes T_\eta)v, \Lambda(a) \otimes \Gamma(b) \rangle = 0$.

Hence we get that $(S_\omega \otimes T_\eta)v = 0$.

Therefore, the remarks after result 2.2 implies that $v = 0$. ■

So we can give the following definition.

Definition 7.3 We define the closed linear map $\Lambda \otimes \Gamma$ from within $A \otimes B$ into $H \otimes K$ such that $\mathcal{N}_\varphi \odot \mathcal{N}_\psi$ is a core for $\Lambda \otimes \Gamma$ and $(\Lambda \otimes \Gamma)(a \otimes b) = \Lambda(a) \otimes \Gamma(b)$ for every $a \in \mathcal{N}_\varphi$ and $b \in \mathcal{N}_\psi$.

Remark 7.4 • We have for every $x \in A \odot B$ and $y \in \mathcal{N}_\varphi \odot \mathcal{N}_\psi$ that xy belongs to $\mathcal{N}_\varphi \odot \mathcal{N}_\psi$ and $(\Lambda \otimes \Gamma)(xy) = (\pi \otimes \theta)(x) (\Lambda \otimes \Gamma)(y)$. This implies easily that $D(\Lambda \otimes \Gamma)$ is a left ideal in $A \otimes B$ and that $(\Lambda \otimes \Gamma)(xy) = (\pi \otimes \theta)(x) (\Lambda \otimes \Gamma)(y)$ for every $x \in A \otimes B$ and $y \in D(\Lambda \otimes \Gamma)$.

- Choose $t \in \mathbb{R}$. Then it is easy to check for every $x \in \mathcal{N}_\varphi \odot \mathcal{N}_\psi$ that $(\sigma \otimes \tau)_t(x)$ belongs to $\mathcal{N}_\varphi \odot \mathcal{N}_\psi$ and $(\Lambda \otimes \Gamma)((\sigma \otimes \tau)_t(x)) = (\nabla^{it} \otimes \Delta^{it})(\Lambda \otimes \Gamma)(x)$.

This will imply for every $x \in D(\Lambda \otimes \Gamma)$ that $(\sigma \otimes \tau)_t(x)$ belongs to $D(\Lambda \otimes \Gamma)$ and $(\Lambda \otimes \Gamma)((\sigma \otimes \tau)_t(x)) = (\nabla^{it} \otimes \Delta^{it})(\Lambda \otimes \Gamma)(x)$.

In the beginning of section 6, we introduced the sub- $*$ -algebras C_φ of $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$ and C_ψ of $\mathcal{N}_\psi \cap \mathcal{N}_\psi^*$. We have also that $C_\varphi \subseteq D(\sigma_z)$ and $\sigma_z(C_\varphi) \subseteq C_\varphi$ and that $C_\psi \subseteq D(\tau_z)$ and $\tau_z(C_\psi) \subseteq C_\psi$ for every $z \in \mathbb{C}$. We know also that C_φ is a core for Λ and that C_ψ is a core for Γ .

Define the $*$ -algebra $C = C_\varphi \odot C_\psi$. Then C is a core for $\Lambda \otimes \Gamma$.

By the remarks after definition 7.1, we see that easily that C consists of analytic elements with respect to $(\sigma \otimes \tau)_z$ and that $(\sigma \otimes \tau)_z(C) \subseteq C$ for every $z \in \mathbb{C}$.

So we get in particular that $C \subseteq D((\sigma \otimes \tau)_{\frac{i}{2}})$ and that $(\sigma \otimes \tau)_{\frac{i}{2}}(C)^* \subseteq C \subseteq D(\Lambda \otimes \Gamma)$.

Using the remarks after definition 7.1 and the fact that $\varphi(b^*a) = \varphi(\sigma_{\frac{i}{2}}(a)\sigma_{\frac{i}{2}}(b)^*)$ for every $a, b \in \mathcal{N}_\varphi \cap D(\sigma_{\frac{i}{2}})$ and $\psi(b^*a) = \psi(\tau_{\frac{i}{2}}(a)\tau_{\frac{i}{2}}(b)^*)$ for every $a, b \in \mathcal{N}_\psi \cap D(\tau_{\frac{i}{2}})$, it is not difficult to check that

$$\|(\Lambda \otimes \Gamma)(x)\|^2 = \|(\Lambda \otimes \Gamma)((\sigma \otimes \tau)_{\frac{i}{2}}(x)^*)\|^2$$

for every $x \in C$.

So we see that the ingredients $A \otimes B$, $H \otimes K$, $\Lambda \otimes \Gamma$, $\pi \otimes \theta$, $\sigma \otimes \theta$, $\nabla \otimes \Delta$ satisfy the conditions of the first construction procedure of section 5. Therefore we can use definition 5.11.

Definition 7.5 We define the weight $\varphi \otimes \psi$ on $A \otimes B$ such that $(H \otimes K, \Lambda \otimes \Gamma, \pi \otimes \theta)$ is a GNS-construction for $\varphi \otimes \psi$.

So we have in particular that $\mathcal{N}_{\varphi \otimes \psi} = D(\Lambda \otimes \Gamma)$, so $\mathcal{N}_\varphi \odot \mathcal{N}_\psi \subseteq \mathcal{N}_{\varphi \otimes \psi}$.

Proposition 5.6 implies the following one :

Proposition 7.6 The weight $\varphi \otimes \psi$ is a KMS-weight with modular group $\sigma \otimes \tau$.

Concerning the modular conjugation we have the following obvious result :

Proposition 7.7 With respect to the GNS-construction, the modular conjugation of $\varphi \otimes \psi$ is given by $J \otimes I$, the modular operator of $\varphi \otimes \psi$ is given by $\nabla \otimes \Delta$.

Call J' the modular conjugation of $\varphi \otimes \psi$. Using definition 6.3 and the remarks after definition 7.1, it is easy to check that $J'(\Lambda(a) \otimes \Gamma(b)) = J\Lambda(a) \otimes I\Gamma(b)$ for every $a \in C_\varphi$ and $b \in C_\psi$. This implies immediately that $J' = J \otimes I$. The result about the modular operator follow from the remarks after definition 7.3.

Definition 7.3 also implies easily the following result :

Result 7.8 Consider $a \in \mathcal{M}_\varphi$ and $b \in \mathcal{M}_\psi$. Then $a \otimes b$ belongs to $\mathcal{M}_{\varphi \otimes \psi}$ and $(\varphi \otimes \psi)(a \otimes b) = \varphi(a) \psi(b)$.

There is also another characterization of $\varphi \otimes \psi$.

Result 7.9 We have the following properties :

1. We have for every $\omega \in \mathcal{F}_\varphi$ and $\eta \in \mathcal{F}_\psi$ that $\omega \otimes \eta$ belongs to $\mathcal{F}_{\varphi \otimes \psi}$.
2. We have for every $\omega \in \mathcal{G}_\varphi$ and $\eta \in \mathcal{G}_\psi$ that $\omega \otimes \eta$ belongs to $\mathcal{G}_{\varphi \otimes \psi}$.

Proof : Consider $\omega \in \mathcal{F}_\varphi$ and $\eta \in \mathcal{F}_\psi$.

Define the operator $S_\omega \in B(H)$ such that $\langle S_\omega \Lambda(a), \Lambda(b) \rangle = \omega(b^* a)$ for every $a, b \in \mathcal{N}_\varphi$.

Define also the operator $T_\eta \in B(K)$ such that $\langle T_\eta \Gamma(a), \Gamma(b) \rangle = \eta(b^* a)$ for every $a, b \in \mathcal{N}_\psi$.

Then it easy to see that

$$\langle (S_\omega \otimes T_\eta)(\Lambda \otimes \Gamma)(x), (\Lambda \otimes \Gamma)(y) \rangle = (\omega \otimes \eta)(y^* x)$$

for every $x, y \in \mathcal{N}_\varphi \odot \mathcal{N}_\psi$. As usual, this implies that

$$\langle (S_\omega \otimes T_\eta)(\Lambda \otimes \Gamma)(x), (\Lambda \otimes \Gamma)(y) \rangle = (\omega \otimes \eta)(y^* x)$$

for every $x, y \in \mathcal{N}_{\varphi \otimes \psi}$.

Because $0 \leq S_\omega \leq 1$ and $0 \leq T_\eta \leq 1$, this implies immediately that $\omega \otimes \eta$ belongs to $\mathcal{F}_{\varphi \otimes \psi}$.

The second result follows from the first one. ■

Proposition 7.10 Consider $x \in (A \otimes B)^+$. Then

$$(\varphi \otimes \psi)(x) = \sup \{ (\omega \otimes \eta)(x) \mid \omega \in \mathcal{F}_\varphi, \eta \in \mathcal{F}_\psi \} .$$

Proof : By the previous proposition and the lower semi-continuity of $\varphi \otimes \psi$, we get immediately the inequality

$$(\varphi \otimes \psi)(x) \geq \sup \{ (\omega \otimes \eta)(x) \mid \omega \in \mathcal{F}_\varphi, \eta \in \mathcal{F}_\psi \} .$$

By proposition 6.5, we have truncating sequences $(u_k)_{k \in K}$ for φ and $(v_l)_{l \in L}$ for ψ .

We define for every $k \in K$ the operator $S_k \in B(H)$ such that $S_k \Lambda(a) = \Lambda(a u_k)$ for every $a \in \mathcal{N}_\varphi$.

In a similar way, we define for every $l \in L$ the operator $T_l \in B(K)$ such that $T_l \Gamma(b) = \Gamma(b v_l)$ for every $b \in \mathcal{N}_\psi$.

Then we get immediately that

- We have that $u_k \otimes v_l$ belongs to $\mathcal{N}_{\varphi \otimes \psi}$ for every $k \in K, l \in L$.
- $\|S_k \otimes T_l\| \leq 1$ for every $k \in K$ and $l \in L$.
- The net $(u_k \otimes v_l)_{(k,l) \in K \times L}$ converges strictly to 1.
- The net $(S_k \otimes T_l)_{(k,l) \in K \times L}$ converges strongly to 1.

Choose $k \in K, l \in L$.

Then it is easy to see that $(S_k \otimes T_l)(\Lambda \otimes \Gamma)(y) = (\Lambda \otimes \Gamma)(y(u_k \otimes v_l))$ for every $y \in \mathcal{N}_\varphi \odot \mathcal{N}_\psi$.

As usual, this implies that $(S_k \otimes T_l)(\Lambda \otimes \Gamma)(y) = (\Lambda \otimes \Gamma)(y(u_k \otimes v_l))$ for every $y \in \mathcal{N}_{\varphi \otimes \psi}$.

So we see that the net $(u_k \otimes v_l)_{(k,l) \in K \otimes L}$ satisfies the conditions of proposition 3.8.

For every $k \in K$, we define $\omega_k \in \mathcal{F}_\varphi$ such that $\omega_k(a) = \varphi(u_k^* a u_k)$ for every $a \in A$.

For every $l \in L$, we define $\eta_l \in \mathcal{F}_\psi$ such that $\eta_l(b) = \varphi(v_l^* b v_l)$ for every $b \in B$.

It is clear that $(\omega_k \otimes \eta_l)(y) = (\varphi \otimes \eta)((u_k \otimes v_l)^* y (u_k \otimes v_l))$ for every $y \in A \otimes B$.

Therefore, proposition 3.8 implies that

$$(\varphi \otimes \psi)(x) = \sup \{ (\omega_k \otimes \eta_l)(x) \mid k \in K, l \in L \} \leq \sup \{ (\omega \otimes \eta)(x) \mid \omega \in \mathcal{F}_\varphi, \eta \in \mathcal{F}_\psi \}.$$

■

Corollary 7.11 *We have the following properties :*

1. *We have for every $x \in (A \otimes B)^+$ that the net $((\omega \otimes \eta)(x))_{(\omega, \eta) \in \mathcal{G}_\varphi \times \mathcal{G}_\psi}$ converges to $(\varphi \otimes \psi)(x)$.*
2. *We have for every $x \in \mathcal{M}_{\varphi \otimes \psi}$ that the net $((\omega \otimes \eta)(x))_{(\omega, \eta) \in \mathcal{G}_\varphi \times \mathcal{G}_\psi}$ converges to $(\varphi \otimes \psi)(x)$.*

8 Absolutely continuous KMS-weights

Consider a C^* -algebra A and a KMS-weight φ on A with modular group σ .

Let δ be a strictly positive element affiliated with A such that there exists a strictly positive number λ such that $\sigma_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$.

In this section we are going to use one of the construction procedures to define the KMS-weight $\varphi(\delta^{\frac{1}{2}} \cdot \delta^{\frac{1}{2}})$ and prove some useful properties about this weight.

It is not possible to use the definition of [14], because it is assumed in this paper that $\sigma_t(\delta) = \delta$ for every $t \in \mathbb{R}$. Our construction procedures from section 5 allows us to go about it in another way.

First, we introduce some notations. Take a GNS-construction (H, Λ, π) for φ . We denote the modular conjugation of φ in this GNS-construction by J and the modular operator by ∇ .

We will extensively have to truncate δ in order to make things bounded. We will also need these truncations to behave well with respect to σ . For this, we will use the following elements in $M(A)$.

Notation 8.1 *Consider $n \in \mathbb{N}$. Then we define $e_n \in M(A)$ such that*

$$e_n a = \frac{n}{\sqrt{\pi}} \int \exp(-n^2 t^2) \delta^{it} a dt$$

for every $a \in A$.

The behaviour of these elements is described in the next proposition.

Proposition 8.2 *We have the following properties.*

1. *The sequence $(e_n)_{n=1}^\infty$ is bounded by 1 and converges strictly to 1.*
2. *We have for every $n \in \mathbb{N}$ that e_n is strictly analytic with respect to σ .*
3. *Let $y \in \mathbb{C}$. Then the sequence $(\sigma_y(e_n))_{n=1}^\infty$ is bounded and converges strictly to 1.*

4. Consider $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then e_n is a left and right multiplier of δ^z and $\delta^z e_n = e_n \delta^z$.
5. Let $n \in \mathbb{N}$ and $y, z \in \mathbb{C}$. Then $\delta^z e_n$ is a left and right multiplier of δ^y and $(\delta^z e_n) \delta^y = \delta^y (\delta^z e_n) = \delta^{y+z} e_n$.
6. Consider $n \in \mathbb{N}$ and $y, z \in \mathbb{C}$. Then $\sigma_y(e_n)$ is a left and right multiplier of δ^z and $\sigma_y(e_n) \delta^z = \delta^z \sigma_y(e_n)$.
7. Consider $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then $\delta^z e_n$ is strictly analytic with respect to σ and $\sigma_y(\delta^z e_n) = \lambda^{yz} \delta^z \sigma_y(e_n)$ for every $y \in \mathbb{C}$.

The results of this proposition will be frequently used in the rest of this section without reference to this proposition. The proof of it is a standard exercise in the use of analytic continuations and will be left out.

We define the set

$$N_0 = \{ a \in A \mid a \text{ is a left multiplier of } \delta^{\frac{1}{2}} \text{ and } a \delta^{\frac{1}{2}} \text{ belongs to } \mathcal{N}_\varphi \} .$$

Introducing this set is of course a very natural thing to do in this case.

It is clear that this set is a left ideal in $M(A)$.

Lemma 8.3 *The mapping $N_0 \rightarrow H : a \mapsto \Lambda(a \delta^{\frac{1}{2}})$ is closable*

Proof: Choose a sequence $(x_k)_{k=1}^\infty$ in N_0 and $v \in H$ such that $(x_k)_{k=1}^\infty$ converges to 0 and $(\Lambda(x_k \delta^{\frac{1}{2}}))_{k=1}^\infty$ converges to v . Choose $n \in \mathbb{N}$.

Take $l \in \mathbb{N}$.

Because e_n is a left multiplier of $\delta^{\frac{1}{2}}$, we have that $x_l e_n$ is a left multiplier of $\delta^{\frac{1}{2}}$ and

$$(x_l e_n) \delta^{\frac{1}{2}} = x_l (e_n \delta^{\frac{1}{2}}) = x_l (\delta^{\frac{1}{2}} e_n) . \quad (\text{a})$$

We know that x_l is a left multiplier of $\delta^{\frac{1}{2}}$, so this last equality implies also that $(x_l e_n) \delta^{\frac{1}{2}} = (x_l \delta^{\frac{1}{2}}) e_n$. Because $x_l \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and e_n belongs to $D(\overline{\sigma}_{\frac{i}{2}})$, this equality and proposition 6.4 imply that $(x_l e_n) \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda((x_l e_n) \delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(x_l \delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Gamma(x_l) . \quad (\text{b})$$

Equality (a) implies that $((x_k e_n) \delta^{\frac{1}{2}})_{k=1}^\infty$ converges to 0, whereas equality (b) implies that $(\Lambda((x_k e_n) \delta^{\frac{1}{2}}))_{k=1}^\infty$ converges to $J\pi(\sigma_{\frac{i}{2}}(e_n))^* Jv$.

Therefore, the closedness of Λ implies that $J\pi(\sigma_{\frac{i}{2}}(e_n))^* Jv = 0$

Because $(\pi(\sigma_{\frac{i}{2}}(e_n)))_{n=1}^\infty$ converges strongly* to 1, this last equation implies that $v = 0$. ■

Therefore, we can give the following definition.

Definition 8.4 *We define Γ as the closure of the linear mapping $N_0 \rightarrow H : a \mapsto \Lambda(a \delta^{\frac{1}{2}})$. The domain of Γ will be denoted by N .*

So Γ is a closed linear mapping from N into H such that N_0 is a core for Γ and $\Gamma(a) = \Lambda(a \delta^{\frac{1}{2}})$ for every $a \in N_0$. It is also easy to check that N is a left ideal in $M(A)$ and that $\Gamma(xa) = \pi(x)\Gamma(a)$ for every $x \in M(A)$ and $a \in N$.

Lemma 8.5 Consider $a \in \mathcal{N}_\varphi$, $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then $a(\delta^z e_n)$ belongs to N_0 and $\Gamma(a(\delta^z e_n)) = J\pi(\sigma_{\frac{i}{2}}(\delta^{z+\frac{1}{2}} e_n))^* J\Lambda(a)$.

Proof : We know that $\delta^z e_n$ is a left multiplier of $\delta^{\frac{1}{2}}$ and $(\delta^z e_n) \delta^{\frac{1}{2}} = \delta^{z+\frac{1}{2}} e_n$. This implies that $a(\delta^z e_n)$ is a left multiplier of $\delta^{\frac{1}{2}}$ and $(a(\delta^z e_n)) \delta^{\frac{1}{2}} = a(\delta^{z+\frac{1}{2}} e_n)$. Because a belongs to $\mathcal{N}_\varphi$ and $\delta^{z+\frac{1}{2}} e_n$ belongs to $D(\overline{\sigma}_{\frac{i}{2}})$, proposition 6.10 implies that $(a(\delta^z e_n)) \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and $\Lambda((a(\delta^z e_n)) \delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(\delta^{z+\frac{1}{2}} e_n))^* J\Lambda(a)$. This gives by definition that $a(\delta^z e_n)$ belongs to N_0 and $\Gamma(a(\delta^z e_n)) = J\pi(\sigma_{\frac{i}{2}}(\delta^{z+\frac{1}{2}} e_n))^* J\Lambda(a)$. ■

Result 8.6 The set N is dense in A and the set $\Gamma(N)$ is dense in H .

Proof : Choose $a \in \mathcal{N}_\varphi$. The previous lemma implies for every $n \in \mathbb{N}$ that $a e_n$ belongs to N_0 . Because $(e_n)_{n=1}^\infty$ converges strictly to 1, we see that $(a e_n)_{n=1}^\infty$ converges to a . Hence, the density of $\mathcal{N}_\varphi$ in A implies that N_0 is dense in A .

Choose $a \in \mathcal{N}_\varphi$. The previous lemma implies for every $n \in \mathbb{N}$ that $a(\delta^{-\frac{1}{2}} e_n)$ belongs to N_0 and $\Gamma(a(\delta^{-\frac{1}{2}} e_n)) = J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(a)$. Because $(\pi(\sigma_{\frac{i}{2}}(e_n)))_{n=1}^\infty$ converges strongly* to 1. This implies that $(\Gamma(a(\delta^{-\frac{1}{2}} e_n)))_{n=1}^\infty$ converges to $\Lambda(a)$. Hence, the density of $\Lambda(\mathcal{N}_\varphi)$ in H implies that $\Gamma(N_0)$ is dense in H . ■

So we have objects A, H, N, Λ, π on which we want to apply the second procedure of section 5. In order to do so, we have to introduce the KMS-characteristics.

Definition 8.7 We define the norm continuous one-parameter group σ' on A such that $\sigma'_t(a) = \delta^{it} \sigma_t(a) \delta^{-it}$ for every $t \in \mathbb{R}$ and $a \in A$.

It is not difficult to see that $\sigma'_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$.

Result 6.13 implies the following lemma.

Lemma 8.8 We have for every $s, t \in \mathbb{R}$ that $J\pi(\delta)^{it} J$ and $\pi(\delta)^{is}$ commute.

The following result is true because $\pi(\sigma_s(a)) = \nabla^{is} \pi(a) \nabla^{-is}$ for every $a \in A$, $s \in \mathbb{R}$ and the fact that $\sigma_s(\delta^{it}) = \lambda^{ist} \delta^{it}$ for $s, t \in \mathbb{R}$ (which is true by assumption).

Lemma 8.9 Consider $s, t \in \mathbb{R}$. Then $\nabla^{is} \pi(\delta)^{it} = \lambda^{ist} \pi(\delta)^{it} \nabla^{is}$

This will imply the following lemma.

Lemma 8.10 We have for every $s, t \in \mathbb{R}$ that $J\pi(\delta)^{it} J \pi(\delta)^{it}$ and ∇^{is} commute.

Proof : We have that

$$\begin{aligned} J\pi(\delta)^{it} J \pi(\delta)^{it} \nabla^{is} &= \lambda^{-ist} J\pi(\delta)^{it} J \nabla^{is} \pi(\delta)^{it} = \lambda^{-ist} J\pi(\delta)^{it} \nabla^{is} J \pi(\delta)^{it} \\ &\stackrel{(*)}{=} \lambda^{-ist} \lambda^{ist} J \nabla^{is} \pi(\delta)^{it} J \pi(\delta)^{it} = \nabla^{is} J\pi(\delta)^{it} J \pi(\delta)^{it} . \end{aligned}$$

In (*), we used that J is antilinear. ■

Lemmas 8.8 and 8.10 imply that the mapping $\mathbb{R} \rightarrow B(H) : t \mapsto J\pi(\delta)^{it} J \pi(\delta)^{it} \nabla^{it}$ is a strongly continuous unitary group representation on H . By the Stone theorem, this justifies the following definition.

Definition 8.11 We define the positive injective operator ∇ (pronounced 'nabla prime') in H such that $\nabla^{it} = J\pi(\delta)^{it}J\pi(\delta)^{it}\nabla^{it}$ for every $t \in \mathbb{R}$.

Proposition 8.12 Consider $a \in \mathcal{N}_\varphi$ and $t \in \mathbb{R}$. Then $\sigma'_t(a)$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(\sigma'_t(a)) = \lambda^{\frac{t}{2}} \nabla^{it} \Lambda(a)$.

Proof : We know by definition that $\sigma_t(a)$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(\sigma_t(a)) = \nabla^{it} \Lambda(a)$. This implies that $\delta^{it} \sigma_t(a)$ belongs to $\mathcal{N}_\varphi$ and that $\Lambda(\delta^{it} \sigma_t(a)) = \pi(\delta)^{it} \Lambda(\sigma_t(a)) = \pi(\delta)^{it} \nabla^{it} \Lambda(a)$. Therefore, proposition 6.30 implies that $\delta^{it} \sigma_t(a) \delta^{-it}$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(\delta^{it} \sigma_t(a) \delta^{-it}) = \lambda^{\frac{t}{2}} J\pi(\delta)^{it} J \Lambda(\delta^{it} \sigma_t(a)) = \lambda^{\frac{t}{2}} J\pi(\delta)^{it} J \pi(\delta)^{it} \nabla^{it} \Lambda(a) .$$

Looking at the definitions of σ'_t and ∇ , the lemma follows. ■

Proposition 8.13 Consider $a \in N$ and $t \in \mathbb{R}$. Then $\sigma_t(a)$ belongs to N and $\Gamma(\sigma_t(a)) = \lambda^{-\frac{t}{2}} \nabla^{it} \Gamma(a)$.

Proof : Choose $b \in N_0$.

Because b is a left multiplier of $\delta^{\frac{1}{2}}$ and $\sigma_t(\delta) = \lambda^t \delta$, we get that $\sigma_t(b)$ is a left multiplier of $\delta^{\frac{1}{2}}$ and

$$\sigma_t(b) \delta^{\frac{1}{2}} = \lambda^{-\frac{t}{2}} \sigma_t(b \delta^{\frac{1}{2}}) .$$

Because $b \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$, this implies that $\sigma_t(b) \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(\sigma_t(b) \delta^{\frac{1}{2}}) = \lambda^{-\frac{t}{2}} \Lambda(\sigma_t(b \delta^{\frac{1}{2}})) = \lambda^{-\frac{t}{2}} \nabla^{it} \Lambda(b \delta^{\frac{1}{2}}) = \lambda^{-\frac{t}{2}} \nabla^{it} \Gamma(b) .$$

So we see that $\sigma_t(b)$ belongs to N_0 and

$$\Gamma(\sigma_t(b)) = \Lambda(\sigma_t(b) \delta^{\frac{1}{2}}) = \lambda^{-\frac{t}{2}} \nabla^{it} \Gamma(b) .$$

The result follows easily because N_0 is a core for Γ . ■

Proposition 8.14 Consider $a \in N$ and $t \in \mathbb{R}$. Then $\sigma'_t(a)$ belongs to N and $\Gamma(\sigma'_t(a)) = \nabla^{it} \Gamma(a)$.

Proof : Choose $b \in N_0$.

Because b is a left multiplier of $\delta^{\frac{1}{2}}$ and $\sigma'_t(\delta) = \lambda^t \delta$, we get that $\sigma'_t(b)$ is a left multiplier of $\delta^{\frac{1}{2}}$ and

$$\sigma'_t(b) \delta^{\frac{1}{2}} = \lambda^{-\frac{t}{2}} \sigma'_t(b \delta^{\frac{1}{2}}) .$$

Because $b \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$, this implies that $\sigma'_t(b) \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(\sigma'_t(b) \delta^{\frac{1}{2}}) = \lambda^{-\frac{t}{2}} \Lambda(\sigma'_t(b \delta^{\frac{1}{2}})) \stackrel{(*)}{=} \lambda^{-\frac{t}{2}} \lambda^{\frac{t}{2}} \nabla^{it} \Lambda(b \delta^{\frac{1}{2}}) = \nabla^{it} \Gamma(b) ,$$

where we used proposition 8.12 in equality (*). The results follows easily because N_0 is a core for Γ . ■

Proposition 8.15 Consider $x \in N$ and $a \in D(\sigma'_{\frac{1}{2}})$. Then xa belongs to N and

$$\Gamma(xa) = J\pi(\sigma'_{\frac{1}{2}}(a))^* J \Gamma(x) .$$

Proof : Define the norm continuous one-parameter group τ on A such that $\tau_t(a) = \delta^{it}a\delta^{-it}$ for every $a \in A$ and $t \in \mathbb{R}$. Then we have that $\sigma_s\tau_t = \tau_t\sigma_s$ for every $s, t \in \mathbb{R}$ and $\sigma'_t(a) = \tau_t(\sigma'_t(a))$ for every $t \in \mathbb{R}$. This implies that $\sigma_{\frac{i}{2}}\tau_{\frac{i}{2}}$ is closable and that the closure is equal to $\sigma'_{\frac{i}{2}}$ (see [9]).

Take $y \in N_0$. So y is a left multiplier of $\delta^{\frac{1}{2}}$ and $y\delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$.

Choose $b \in D(\sigma_{\frac{i}{2}}\tau_{\frac{i}{2}})$.

Because b belongs to $D(\tau_{\frac{i}{2}})$, the element b is a middle multiplier of $\delta^{-\frac{1}{2}}$, $\delta^{\frac{1}{2}}$ and $\delta^{-\frac{1}{2}}b\delta^{\frac{1}{2}} = \tau_{\frac{i}{2}}(b)$.

We have by assumption that y is a left multiplier of $\delta^{\frac{1}{2}}$. This implies that $y\delta^{\frac{1}{2}}$ is a left multiplier of $\delta^{-\frac{1}{2}}$ and $(y\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}} = y$.

This in turn implies that $((y\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}})b$ is a left multiplier of $\delta^{\frac{1}{2}}$ and

$$[(y\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}}]b\delta^{\frac{1}{2}} = (y\delta^{\frac{1}{2}})(\delta^{-\frac{1}{2}}b\delta^{\frac{1}{2}}) = (y\delta^{\frac{1}{2}})\tau_{\frac{i}{2}}(b).$$

Combining this with the fact that $((y\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}})b = yb$, we see that yb is a left multiplier of $\delta^{\frac{1}{2}}$ and $(yb)\delta^{\frac{1}{2}} = (y\delta^{\frac{1}{2}})\tau_{\frac{i}{2}}(b)$.

Because $y\delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and $\tau_{\frac{i}{2}}(b)$ belongs to $D(\sigma_{\frac{i}{2}})$, this last equation implies that $(yb)\delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda((yb)\delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(\tau_{\frac{i}{2}}(b)))^*J\Lambda(y\delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(\tau_{\frac{i}{2}}(b)))^*J\Gamma(y).$$

Therefore, yb belongs to N_0 and $\Gamma(yb) = J\pi(\sigma_{\frac{i}{2}}(\tau_{\frac{i}{2}}(b)))^*J\Gamma(y)$

So we see that $\Gamma(yb) = J\pi(\sigma'_{\frac{i}{2}}(b))^*J\Gamma(y)$. (*)

Because $D(\sigma_{\frac{i}{2}}\tau_{\frac{i}{2}})$ is a core for $\sigma'_{\frac{i}{2}}$, there exists a sequence $(a_n)_{n=1}^\infty$ in $D(\sigma_{\frac{i}{2}}\tau_{\frac{i}{2}})$ such that $(a_n)_{n=1}^\infty$ converges to a and $(\sigma'_{\frac{i}{2}}(a_n))_{n=1}^\infty$ converges to $\sigma'_{\frac{i}{2}}(a)$.

By result (*) we have for every $n \in \mathbb{N}$ that ya_n belongs to N_0 and

$$\Gamma(ya_n) = J\pi(\sigma'_{\frac{i}{2}}(a_n))^*J\Gamma(y)$$

From this, we conclude that the sequence $(\Gamma(ya_n))_{n=1}^\infty$ converges to $J\pi(\sigma'_{\frac{i}{2}}(a))^*J\Gamma(y)$. It is also clear that $(ya_n)_{n=1}^\infty$ converges to ya so the closedness of Γ implies that ya belongs to N and $\Gamma(ya) = J^*\pi(\sigma'_{\frac{i}{2}}(a))^*J\Gamma(y)$.

The proposition follows because N_0 is a core for Γ . ■

At last we are in a position to define $\varphi(\delta^{\frac{1}{2}} \cdot \delta^{\frac{1}{2}})$.

Definition 8.16 *We have proven in the previous part of the section that the ingredients $A, H, N, \Lambda, \pi, \sigma', \nabla$ and J satisfy the conditions of the second construction procedure of section 5. The weight which is obtained from this ingredients is denoted by φ_δ (see definition 5.12).*

We want to forget about this definition, but will instead repeat the determining properties. Previously, we have defined a norm-continuous one-parameter group σ' on A such that $\sigma'_t(a) = \delta^{it}\sigma_t(a)\delta^{-it}$ for every $t \in \mathbb{R}$ and $a \in A$. Then

Theorem 8.17 *We have that φ_δ is a KMS-weight with modular group σ' .*

The weight φ_δ is completely determined by the following proposition.

Theorem 8.18 *We have that (H, Γ, π) is a GNS-construction for φ_δ*

Remember that the set

$$N_0 = \{ a \in A \mid a \text{ is a left multiplier of } \delta^{\frac{1}{2}} \text{ and } a \delta^{\frac{1}{2}} \text{ belongs to } \mathcal{N}_\varphi \} .$$

is a core for Γ and that $\Gamma(a) = \Lambda(a \delta^{\frac{1}{2}})$ for every $a \in N_0$.

Repeating the results of lemmas 8.8 and 8.10 gives

- We have for every $s, t \in \mathbb{R}$ that $J\pi(\delta)^{it}J$ and $\pi(\delta)^{is}$ commute.
- We have for every $s, t \in \mathbb{R}$ that $J\pi(\delta)^{it}J\pi(\delta)^{it}$ and ∇^{is} commute.

Also, the injective positive operator $\mathbf{V}$ in H was defined in such a way that $\mathbf{V}^{it} = J\pi(\delta)^{it}J\pi(\delta)^{it}\nabla^{it}$ for every $t \in \mathbb{R}$. Then :

Proposition 8.19 *We have that $\mathbf{V}$ is the modular operator for φ_δ in the GNS-construction (H, Γ, π) .*

Later, we will prove that the modular conjugation for φ_δ is a scalar multiple of J .

The following relative invariance properties are also valid.

- Consider $a \in \mathcal{N}_\varphi$ and $t \in \mathbb{R}$. Then $\sigma'_t(a)$ belongs to $\mathcal{N}_\varphi$ and $\Lambda(\sigma'_t(a)) = \lambda^{\frac{t}{2}} \mathbf{V}^{it} \Lambda(a)$.
- Consider $a \in \mathcal{N}_{\varphi_\delta}$ and $t \in \mathbb{R}$. Then $\sigma_t(a)$ belongs to $\mathcal{N}_{\varphi_\delta}$ and $\Gamma(\sigma_t(a)) = \lambda^{-\frac{t}{2}} \nabla^{it} \Gamma(a)$.

These imply the following proposition.

Proposition 8.20 *Consider $t \in \mathbb{R}$. Then $\varphi \sigma'_t = \lambda^t \varphi$ and $\varphi_\delta \sigma_t = \lambda^{-t} \varphi_\delta$.*

Because (H, Γ, π) is a GNS-construction for φ_δ , proposition 6.22 implies the following result.

Proposition 8.21 *The weight φ_δ is faithful $\Leftrightarrow \varphi$ is faithful*

Because $\sigma'_t(\delta) = \lambda^t \delta$ for every $t \in \mathbb{R}$, also the following proposition will be true (see proposition 8.2).

Proposition 8.22 *We have the following properties.*

1. *We have for every $n \in \mathbb{N}$ that e_n is strictly analytic with respect to σ' .*
2. *Let $y \in \mathbb{C}$. Then the sequence $(\sigma'_y(e_n))_{n=1}^\infty$ is bounded and converges strictly to 1.*
3. *Consider $n \in \mathbb{N}$ and $y, z \in \mathbb{C}$. Then $\sigma'_y(e_n)$ is a left and right multiplier of δ^z and $\sigma'_y(e_n) \delta^z = \delta^z \sigma'_y(e_n)$.*
4. *Consider $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then $\delta^z e_n$ is strictly analytic with respect to σ' and $\sigma'_y(\delta^z e_n) = \lambda^{yz} \delta^z \sigma'_y(e_n)$ for every $y \in \mathbb{C}$.*

In the next part of this section, we will prove a formula which formally says that $\varphi_\delta(x) = \varphi(\delta^{\frac{1}{2}} x \delta^{\frac{1}{2}})$ (and something more general).

Lemma 8.23 *We have the following properties :*

- *We have for every $a \in \mathcal{N}_\varphi$, $n \in \mathbb{N}$ and $z \in \mathbb{C}$ that $a(\delta^z e_n)$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta}$.*

- We have for every $x \in \mathcal{M}_\varphi$, $m, n \in \mathbb{N}$ and $y, z \in \mathbb{C}$ that $(\delta^y e_m) x (\delta^z e_n)$ belongs to $\mathcal{M}_\varphi \cap \mathcal{M}_{\varphi_\delta}$.

Proof :

- From lemma 8.5, we know already that $a (\delta^z e_n)$ belongs to $\mathcal{N}_{\varphi_\delta}$. Because a belongs to $\mathcal{N}_\varphi$ and $\delta^z e_n$ belongs to $D(\sigma_{\frac{i}{2}})$, proposition 6.10 implies also that $a (\delta^z e_n)$ belongs to $\mathcal{N}_\varphi$.
- This follows immediately from the first property. ■

Lemma 8.24 Consider $x \in \mathcal{M}_\varphi$. Then $(\varphi(e_n x e_n))_{n=1}^\infty$ converges to $\varphi(x)$.

Proof : Choose $a, b \in \mathcal{N}_\varphi$. We have for every $n \in \mathbb{N}$ that $a e_n, b e_n$ belong to $\mathcal{N}_\varphi$ and $\Lambda(a e_n) = J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(a)$, $\Lambda(b e_n) = J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(b)$ which implies that

$$\varphi(e_n b^* a e_n) = \langle J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(a), J\pi(\sigma_{\frac{i}{2}}(e_n))^* J\Lambda(b) \rangle .$$

Because $(\pi(\sigma_{\frac{i}{2}}(e_n)))_{n=1}^\infty$ converges strongly* to 1, we see that $(\varphi(e_n b^* a e_n))_{n=1}^\infty$ converges to $\langle \Lambda(a), \Lambda(b) \rangle$ which is equal to $\varphi(b^* a)$. ■

Lemma 8.25 Consider $x \in \mathcal{N}_{\varphi_\delta}$, $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then $x (\delta^z e_n)$ belongs to $\mathcal{N}_\varphi$.

Proof : Because $\delta^z e_n$ belongs to $D(\overline{\sigma'_{\frac{i}{2}}})$, we have immediately that $x (\delta^z e_n)$ belongs to $\mathcal{N}_{\varphi_\delta}$. By definition, there exists a sequence $(x_k)_{k=1}^\infty$ in N_0 such that $(x_k)_{k=1}^\infty$ converges to x and $(\Gamma(x_k))_{k=1}^\infty$ converges to $\Gamma(x)$.

Choose $l \in \mathbb{N}$. Because x_l is a left multiplier of $\delta^{\frac{1}{2}}$ and $\delta^{z-\frac{1}{2}} e_n$ is a right multiplier of $\delta^{\frac{1}{2}}$, we have that

$$(x_l \delta^{\frac{1}{2}})(\delta^{z-\frac{1}{2}} e_n) = x_l (\delta^{\frac{1}{2}} (\delta^{z-\frac{1}{2}} e_n)) = x_l (\delta^z e_n) .$$

Because $x_l \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$ and $\delta^{z-\frac{1}{2}} e_n$ belongs to $D(\overline{\sigma'_{\frac{i}{2}}})$, this implies that $x_l (\delta^z e_n)$ belongs to $\mathcal{N}_\varphi$ and

$$\Lambda(x_l (\delta^z e_n)) = J\pi(\sigma_{\frac{i}{2}}(\delta^{z-\frac{1}{2}} e_n))^* J\Lambda(x_l \delta^{\frac{1}{2}}) = J\pi(\sigma_{\frac{i}{2}}(\delta^{z-\frac{1}{2}} e_n))^* J\Gamma(x_l) .$$

From this last equation, we infer that $(\Lambda(x_k (\delta^z e_n)))_{k=1}^\infty$ converges to $J\pi(\sigma_{\frac{i}{2}}(\delta^{z-\frac{1}{2}} e_n))^* J\Gamma(x)$. It is also clear that $(x_k (\delta^z e_n))_{k=1}^\infty$ converges to $x (\delta^z e_n)$. Therefore, the closedness of Λ implies that $x (\delta^z e_n)$ belongs to $\mathcal{N}_\varphi$. ■

Using this lemma, we can even do better.

Lemma 8.26 Consider $x \in \mathcal{N}_{\varphi_\delta}$, $n \in \mathbb{N}$ and $z \in \mathbb{C}$. Then $x (\delta^z e_n)$ belongs to N_0 .

Proof : Because $\delta^z e_n$ is a left multiplier of $\delta^{\frac{1}{2}}$, we get that $x (\delta^z e_n)$ is a left multiplier of $\delta^{\frac{1}{2}}$ and

$$(x (\delta^z e_n)) \delta^{\frac{1}{2}} = x ((\delta^z e_n) \delta^{\frac{1}{2}}) = x (\delta^{z+\frac{1}{2}} e_n)$$

which belongs to $\mathcal{N}_\varphi$ by the previous lemma. The result follows by the definition of N_0 . ■

The following results can be proven in a similar way as for elements in $\mathcal{M}_\varphi$ and $\mathcal{N}_\varphi$.

Lemma 8.27 *We have the following properties.*

- We have for every $a \in \mathcal{N}_{\varphi_\delta}$, $n \in \mathbb{N}$ and $z \in \mathbb{C}$ that $a(\delta^z e_n)$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta}$.
- We have for every $x \in \mathcal{M}_{\varphi_\delta}$, $m, n \in \mathbb{N}$ and $y, z \in \mathbb{C}$ that $(\delta^y e_m)x(\delta^z e_n)$ belongs to $\mathcal{M}_\varphi \cap \mathcal{M}_{\varphi_\delta}$.

Lemma 8.28 *Consider $x \in \mathcal{M}_{\varphi_\delta}$. Then $(\varphi_\delta(e_n x e_n))_{n=1}^\infty$ converges to $\varphi_\delta(x)$.*

We have also the next result.

Corollary 8.29 *The set $\mathcal{N}_\varphi \cap N_0$ is a core for Γ and a core for Λ .*

Proof : Choose $x \in \mathcal{N}_{\varphi_\delta}$. We have by the previous lemmas that $x e_n$ belongs to $\mathcal{N}_\varphi \cap N_0$ for every $n \in \mathbb{N}$. By proposition 6.10, we know that $\Gamma(x e_n) = J' \pi(\sigma'_{\frac{i}{2}}(e_n))^* J' \Gamma(x)$ for every $n \in \mathbb{N}$ (here J' denotes the modular conjugation of φ_δ). This implies that $(\Gamma(x e_n))_{n=1}^\infty$ converges to $\Gamma(x)$. It is also clear that $(x e_n)_{n=1}^\infty$ converges to x .

The statement about Λ is proven in a similar way. ■

Lemma 8.30 *Consider $x \in \mathcal{M}_{\varphi_\delta}$, $z \in \mathbb{C}$ and $n \in \mathbb{N}$. Then $e_n x e_n$ belongs to $\mathcal{M}_{\varphi_\delta}$, $(\delta^{1-z} e_n)x(\delta^z e_n)$ belongs to $\mathcal{M}_\varphi$ and*

$$\varphi_\delta(e_n x e_n) = \lambda^{i(\frac{1}{2}-z)} \varphi((\delta^{1-z} e_n)x(\delta^z e_n)) .$$

Proof : Choose $a, b \in \mathcal{N}_{\varphi_\delta}$. By lemma 8.26, we know that $a e_n, b e_n$ belong to N_0 . This implies that

$$\begin{aligned} \varphi_\delta(e_n b^* a e_n) &= \langle \Gamma(a e_n), \Gamma(b e_n) \rangle = \langle \Lambda((a e_n) \delta^{\frac{1}{2}}), \Lambda((b e_n) \delta^{\frac{1}{2}}) \rangle \\ &= \langle \Lambda(a(\delta^{\frac{1}{2}} e_n)), \Lambda(b(\delta^{\frac{1}{2}} e_n)) \rangle = \varphi((\delta^{\frac{1}{2}} e_n) b^* a (\delta^{\frac{1}{2}} e_n)) . \end{aligned}$$

We know that $\delta^{1-z} e_n$ is a right multiplier of $\delta^{z-\frac{1}{2}}$ and that $\delta^{z-\frac{1}{2}} (\delta^{1-z} e_n) = \delta^{\frac{1}{2}} e_n$. This implies that $(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)$ is a right multiplier of $\delta^{z-\frac{1}{2}}$ and

$$\delta^{z-\frac{1}{2}} [(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)] = (\delta^{\frac{1}{2}} e_n) b^* a (\delta^{\frac{1}{2}} e_n) .$$

So $\delta^{z-\frac{1}{2}} [(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)]$ belongs to $\mathcal{M}_\varphi$ by lemma 8.27.

We know on the other hand that $\delta^{\frac{1}{2}} e_n$ is a left multiplier of $\delta^{z-\frac{1}{2}}$ and that $(\delta^{\frac{1}{2}} e_n) \delta^{z-\frac{1}{2}} = \delta^z e_n$. This implies that $(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)$ is a left multiplier of $\delta^{z-\frac{1}{2}}$ and

$$[(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)] \delta^{z-\frac{1}{2}} = (\delta^{1-z} e_n) b^* a (\delta^z e_n) .$$

So $[(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)] \delta^{z-\frac{1}{2}}$ belongs also to $\mathcal{M}_\varphi$ by 8.27.

Therefore proposition 6.32 implies that

$$\begin{aligned} \varphi_\delta(e_n b^* a e_n) &= \varphi((\delta^{\frac{1}{2}} e_n) b^* a (\delta^{\frac{1}{2}} e_n)) = \varphi(\delta^{z-\frac{1}{2}} [(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)]) \\ &= \lambda^{i(\frac{1}{2}-z)} \varphi([(\delta^{1-z} e_n) b^* a (\delta^{\frac{1}{2}} e_n)] \delta^{z-\frac{1}{2}}) \\ &= \lambda^{i(\frac{1}{2}-z)} \varphi((\delta^{1-z} e_n) b^* a (\delta^z e_n)) . \end{aligned}$$

■

Proposition 8.31 Consider $x \in \mathcal{M}_{\varphi_\delta}$ and $z \in \mathbb{C}$ such that x is a middle multiplier of δ^{1-z} , δ^z and $\delta^{1-z}x\delta^z$ belongs to $\mathcal{M}_\varphi$. Then

$$\varphi_\delta(x) = \lambda^{i(\frac{1}{2}-z)} \varphi(\delta^{1-z}x\delta^z) .$$

Proof : Choose $n \in \mathbb{N}$. By the previous lemma, we know that $e_n x e_n$ belongs to $\mathcal{M}_{\varphi_\delta}$, $(\delta^{1-z}e_n)x(\delta^ze_n)$ belongs to $\mathcal{M}_\varphi$ and

$$\varphi_\delta(e_n x e_n) = \lambda^{i(\frac{1}{2}-z)} \varphi((\delta^{1-z}e_n)x(\delta^ze_n)) .$$

Because e_n is a left multiplier of δ^{1-z} and e_n is a right multiplier of δ^z , we get that

$$e_n(\delta^{1-z}x\delta^z)e_n = (e_n\delta^{1-z})x(\delta^ze_n) = (\delta^{1-z}e_n)x(\delta^ze_n) .$$

So $e_n(\delta^{1-z}x\delta^z)e_n$ belongs to $\mathcal{M}_\varphi$ and

$$\varphi_\delta(e_n x e_n) = \lambda^{i(\frac{1}{2}-z)} \varphi(e_n(\delta^{1-z}x\delta^z)e_n) .$$

The proposition follows by lemmas 8.24 and 8.28. ■

Remark 8.32 We would like to mention the following special cases.

1. Consider $x \in \mathcal{M}_{\varphi_\delta}$ such that x is a middle multiplier of $\delta^{\frac{1}{2}}$, $\delta^{\frac{1}{2}}$ and $\delta^{\frac{1}{2}}x\delta^{\frac{1}{2}}$ belongs to $\mathcal{M}_\varphi$. Then

$$\varphi_\delta(x) = \varphi(\delta^{\frac{1}{2}}x\delta^{\frac{1}{2}}) .$$

2. Consider $x \in \mathcal{M}_{\varphi_\delta}$ such that x is a left multiplier of δ and $x\delta$ belongs to $\mathcal{M}_\varphi$. Then

$$\varphi_\delta(x) = \lambda^{-\frac{i}{2}} \varphi(x\delta) .$$

3. Consider $x \in \mathcal{M}_{\varphi_\delta}$ such that x is a right multiplier of δ and δx belongs to $\mathcal{M}_\varphi$. Then

$$\varphi_\delta(x) = \lambda^{\frac{i}{2}} \varphi(\delta x) .$$

Proposition 8.33 Consider an element $a \in A^+$ and $z \in \mathbb{C}$ such that a is a middle multiplier of δ^{1-z} , δ^z and $\delta^{1-z}a\delta^z$ belongs to $\mathcal{M}_\varphi$. Then a belongs to $\mathcal{M}_{\varphi_\delta}^+$.

Proof : Take $b \in A$ such that $b^*b = a$.

Choose $m \in \mathbb{N}$. Because $\delta^{z-1}e_m$ is a left multiplier of δ^{1-z} and $\delta^{-z}e_m$ is a right multiplier of δ^z , we have that

$$(\delta^{z-1}e_m)(\delta^{1-z}a\delta^z)(\delta^{-z}e_m) = ((\delta^{z-1}e_m)\delta^{1-z})a(\delta^z(\delta^{-z}e_m)) = e_m a e_m = (be_m)^*(be_m) ,$$

Because $\delta^{1-z}a\delta^z$ belongs to $\mathcal{M}_\varphi$, this equality and lemma 8.23 imply that $(be_m)^*(be_m)$ belongs to $\mathcal{M}_{\varphi_\delta}$. Hence be_m belongs to $\mathcal{N}_{\varphi_\delta}$.

Take $k, l \in \mathbb{N}$. By the previous part, we already know that $e_l b^* b e_k$ belongs to $\mathcal{M}_{\varphi_\delta}$.

Because e_l is a right multiplier of δ^{1-z} and e_k is a left multiplier of δ^z , we have that $e_l b^* b e_k$ is a middle multiplier of δ^{1-z} , δ^z and

$$\delta^{1-z}(e_l b^* b e_k)\delta^z = (\delta^{1-z}e_l)b^*b(e_k\delta^z) = (\delta^{1-z}e_l)a(e_k\delta^z)$$

Knowing that e_l is a left multiplier of δ^{1-z} and e_n is a right multiplier of δ^z , we get moreover that

$$e_l (\delta^{1-z} a \delta^z) e_k = (e_l \delta^{1-z}) a (\delta^z e_k) = (\delta^{1-z} e_l) a (e_k \delta^z) = \delta^{1-z} (e_l b^* b e_k) \delta^z .$$

Hence $\delta^{1-z} (e_l b^* b e_k) \delta^z$ belongs to $\mathcal{M}_\varphi$ by lemma 8.23.

Therefore the previous proposition implies that

$$\varphi_\delta(e_l b^* b e_k) = \lambda^{i(\frac{1}{2}-z)} \varphi(\delta^{1-z} (e_l b^* b e_k) \delta^z) = \lambda^{i(\frac{1}{2}-z)} \varphi(e_l (\delta^{1-z} a \delta^z) e_k) .$$

Therefore, we have for all $m, n \in \mathbb{N}$ that

$$\begin{aligned} \|\Gamma(b e_m) - \Gamma(b e_n)\|^2 &= \varphi_\delta(e_m b^* b e_m) - \varphi_\delta(e_m b^* b e_n) - \varphi_\delta(e_n b^* b e_m) + \varphi_\delta(e_n b^* b e_n) \\ &= \lambda^{i(\frac{1}{2}-z)} [\varphi(e_m (\delta^{1-z} a \delta^z) e_m) - \varphi(e_m (\delta^{1-z} a \delta^z) e_n) - \varphi(e_n (\delta^{1-z} a \delta^z) e_m) + \varphi(e_n (\delta^{1-z} a \delta^z) e_n)] \end{aligned}$$

Using this equality and the obvious generalization of lemma 8.24, we see that $(\Gamma(b e_n))_{n=1}^\infty$ is Cauchy and hence convergent. Because we also have that $(b e_n)_{n=1}^\infty$ converges to b , the closedness of Γ imply that b belongs to $\mathcal{N}_{\varphi_\delta}$. Hence, a is an element in $\mathcal{M}_{\varphi_\delta}$. ■

The next proposition is proven in a similar way.

Proposition 8.34 *Consider an element $a \in \mathcal{M}_{\varphi_\delta}$ and $z \in \mathbb{C}$ such that a is a middle multiplier of δ^{1-z} , δ^z and $\delta^{1-z} a \delta^z$ belongs to A^+ . Then $\delta^{1-z} a \delta^z$ belongs to $\mathcal{M}_\varphi^+$.*

Corollary 8.35 *Consider an element $a \in A^+$ such that a is a middle multiplier of $\delta^{\frac{1}{2}}$, $\delta^{\frac{1}{2}}$ and $\delta^{\frac{1}{2}} a \delta^{\frac{1}{2}}$ belongs to A . Then a belongs to $\mathcal{M}_{\varphi_\delta}^+ \Leftrightarrow \delta^{\frac{1}{2}} a \delta^{\frac{1}{2}}$ belongs to $\mathcal{M}_\varphi^+$.*

In the next part of this section, we will prove that the modular conjugation of φ_δ with respect to (H, Γ, π) is a scalar multiple of J .

Notation 8.36 *We define the set $D = \{x \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^* \mid \text{We have for every } z \in \mathbb{C}$*

that

1. x is a right multiplier of δ^z and $\delta^z x$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$
 2. x is a left multiplier of δ^z and $x \delta^z$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$
- } .

It is not difficult to check that D is a sub- $*$ -algebra of $N_0 \cap N_0^* \subseteq \mathcal{N}_{\varphi_\delta} \cap \mathcal{N}_{\varphi_\delta}^*$.

Lemma 8.37 *Consider $x \in \mathcal{N}_{\varphi_\delta} \cap \mathcal{N}_{\varphi_\delta}^*$. Then there exists a sequence $(x_n)_{n=1}^\infty$ in D such that*

1. $(x_n)_{n=1}^\infty$ converges to x
2. $(\Gamma(x_n))_{n=1}^\infty$ converges to $\Gamma(x)$
3. $(\Gamma(x_n^*))_{n=1}^\infty$ converges to $\Gamma(x^*)$

Proof : We define for every $n \in \mathbb{N}$ the element $x_n = e_n x e_n$. Using lemma 8.25, it is not difficult to check that x_n belongs to D for every $n \in \mathbb{N}$. We have also for every $n \in \mathbb{N}$ that

- $\Gamma(x_n) = \Gamma(e_n x e_n) = J \pi(\sigma'_{\frac{1}{2}}(e_n))^* J \pi(e_n) \Gamma(x)$

- $\Gamma(x_n^*) = \Gamma(e_n x^* e_n) = J\pi(\sigma'_{\frac{1}{2}}(e_n))^* J\pi(e_n)\Gamma(x^*)$

This implies that $(x_n)_{n=1}^\infty$ converges to x , $(\Gamma(x_n))_{n=1}^\infty$ converges to $\Gamma(x)$ and $(\Gamma(x_n^*))_{n=1}^\infty$ converges to $\Gamma(x^*)$ $\blacksquare$

It follows immediately that $\Gamma(D)$ is dense in H .

Proposition 8.38 *We have that $\lambda^{\frac{1}{4}} J$ is the modular operator for φ_δ in the GNS-construction (H, Γ, π) .*

Proof : Denote the modular conjugation of φ_δ in the GNS-construction (H, Γ, π) by J' .

Choose $a \in D$.

- We have that a^* is a left multiplier of $\delta^{\frac{1}{2}}$ and that $a^* \delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$. This implies that $\Gamma(a^*) = \Lambda(a^* \delta^{\frac{1}{2}})$ by the definition of Γ . We know moreover that $a \in D(\delta^{\frac{1}{2}})$ and that $\delta^{\frac{1}{2}} a$ belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. This implies that $\Lambda(\delta^{\frac{1}{2}} a)$ belongs to $D(\nabla^{\frac{1}{2}})$ and

$$J\nabla^{\frac{1}{2}}\Lambda(\delta^{\frac{1}{2}}a) = \Lambda((\delta^{\frac{1}{2}}a)^*) = \Lambda(a^*\delta^{\frac{1}{2}}) = \Gamma(a^*) .$$

- Because a belongs to $\mathcal{N}_\varphi$, a belongs to $D(\delta^{\frac{1}{2}})$ and $\delta^{\frac{1}{2}} a$ belongs to $\mathcal{N}_\varphi$, proposition 2.18 implies that $\Lambda(a)$ belongs to $D(\pi(\delta)^{\frac{1}{2}})$ and $\pi(\delta)^{\frac{1}{2}}\Lambda(a) = \Lambda(\delta^{\frac{1}{2}}a)$. So, using the previous part, we get that $\Lambda(a)$ belongs to $D(\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}})$ and

$$J\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}\Lambda(a) = J\nabla^{\frac{1}{2}}\Lambda(\delta^{\frac{1}{2}}a) = \Gamma(a^*) .$$

- We know that a is a left multiplier of $\delta^{\frac{1}{2}}$. This implies that $a\delta^{\frac{1}{2}}$ is a left multiplier of $\delta^{-\frac{1}{2}}$ and that $(a\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}} = a$ which belongs to $\mathcal{N}_\varphi$. Therefore, proposition 6.31 implies that $\Lambda(a\delta^{\frac{1}{2}})$ belongs to $D((J\pi(\delta)J)^{-\frac{1}{2}})$ and

$$(J\pi(\delta)J)^{-\frac{1}{2}}\Lambda(a\delta^{\frac{1}{2}}) = \lambda^{\frac{1}{4}}\Lambda((a\delta^{\frac{1}{2}})\delta^{-\frac{1}{2}}) = \lambda^{\frac{1}{4}}\Lambda(a) .$$

By definition, we have that $\Gamma(a) = \Lambda(a\delta^{\frac{1}{2}})$. Consequently, we have that $\Gamma(a)$ belongs to $D((J\pi(\delta)J)^{-\frac{1}{2}})$ and

$$(J\pi(\delta)J)^{-\frac{1}{2}}\Gamma(a) = \lambda^{\frac{1}{4}}\Lambda(a) .$$

Using the previous part, we see that $\Gamma(a)$ belongs to $D(\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}})$ and

$$J\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}}\Gamma(a) = \lambda^{-\frac{1}{4}}J\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}\Lambda(a) = \lambda^{-\frac{1}{4}}\Gamma(a^*) .$$

The only thing we need from the previous discussion is that $\Gamma(a)$ belongs to $D(\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}})$ and

$$J\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}}\Gamma(a) = \lambda^{-\frac{1}{4}}\Gamma(a^*) . \quad (*)$$

We have that $\pi(\delta)$ and $J\pi(\delta)J$ are strongly commuting (lemma 8.8). Define the positive injective operator S in H such that $S^{it} = \pi(\delta)^{it}(J\pi(\delta)J)^{-it}$ for every $t \in \mathbb{R}$. Therefore, $\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}} \subseteq S^{\frac{1}{2}}$.

We also have that ∇ and S are strongly commuting (lemma 8.10). By definition, we have that $\nabla^{it} = \nabla^{it}S^{it}$ for $t \in \mathbb{R}$. This implies that $\nabla^{\frac{1}{2}}S^{\frac{1}{2}} \subseteq \nabla^{\frac{1}{2}}$.

Combining these two facts, we arrive at the conclusion that $\nabla^{\frac{1}{2}}\pi(\delta)^{\frac{1}{2}}(J\pi(\delta)J)^{-\frac{1}{2}} \subseteq \nabla^{\frac{1}{2}}$.

Using (*), this gives that $\Gamma(a)$ belongs to $D(\nabla^{\frac{1}{2}})$ and $J\nabla^{\frac{1}{2}}\Gamma(a) = \lambda^{-\frac{1}{4}}\Gamma(a^*)$.

On the other hand, the fact that $a \in \mathcal{N}_{\varphi_\delta} \cap \mathcal{N}_{\varphi_\delta}^*$ implies that $J'\nabla^{\frac{1}{2}}\Gamma(a) = \Gamma(a^*)$.

Combining these two equalities results in the equality $\lambda^{\frac{1}{4}}J\Gamma(a^*) = J'\Gamma(a^*)$.

Because $\Gamma(D)$ is dense in H , we get that $J' = \lambda^{\frac{1}{4}}J$. $\blacksquare$

The following two lemmas will be used to prove that δ is uniquely determined in the GNS-representation of φ by the weight φ_δ .

Lemma 8.39 Consider an element $a \in \mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta}$. Then $\Lambda(a)$ belongs to $D(J\pi(\delta^{\frac{1}{2}})J)$ and

$$\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(a) = \Gamma(a) .$$

Consequently, $\|J\pi(\delta^{\frac{1}{2}})J\Lambda(a)\|^2 = \varphi_\delta(a^*a)$.

Proof : Choose $m \in \mathbb{N}$.

Because $a \in \mathcal{N}_\varphi$, we know by lemma 8.23 that ae_m belongs to $\mathcal{N}_\varphi$.

We also know that e_m is a left multiplier of $\delta^{\frac{1}{2}}$, which implies that $(ae_m)\delta^{\frac{1}{2}}$ is a left multiplier of $\delta^{\frac{1}{2}}$ and $(ae_m)\delta^{\frac{1}{2}} = a(e_m\delta^{\frac{1}{2}})$. This last equality ensures that also $(ae_m)\delta^{\frac{1}{2}}$ belongs to $\mathcal{N}_\varphi$.

Hence, proposition 6.31 implies that $\Lambda(ae_m)$ belongs to $D(J\pi(\delta)^{\frac{1}{2}}J)$ and

$$\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(ae_m) = \Lambda((ae_m)\delta^{\frac{1}{2}})$$

So by the definition of Γ , we see that ae_m belongs to $\mathcal{N}_{\varphi_\delta}$ and

$$\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(ae_m) = \Gamma(ae_m)$$

Because $a \in \mathcal{N}_{\varphi_\delta}$ and $e_m \in D(\overline{\sigma}_{\frac{i}{2}}')$, this implies that

$$\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(ae_m) = J\pi(\sigma'_{\frac{i}{2}}(e_m))^* J\Gamma(a)$$

Because $a \in \mathcal{N}_\varphi$ and $e_m \in D(\sigma_{\frac{i}{2}})$, we have also that

$$\Lambda(ae_m) = J\pi(\sigma_{\frac{i}{2}}(e_m))^* J\Lambda(a)$$

So we see that $(\Lambda(ae_n))_{n=1}^\infty$ converges to $\Lambda(a)$ and that $(\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(ae_n))_{n=1}^\infty$ converges to $\Gamma(a)$.

Therefore the closedness of $J\pi(\delta^{\frac{1}{2}})J$ implies that $\Lambda(a)$ belongs to $D(J\pi(\delta^{\frac{1}{2}})J)$ and

$$\lambda^{\frac{i}{4}} J\pi(\delta^{\frac{1}{2}})J\Lambda(a) = \Gamma(a) .$$

■

Lemma 8.40 The set $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta})$ is a core for $J\pi(\delta^{\frac{1}{2}})J$.

Proof : Choose $m \in \mathbb{N}$.

We know already that $\sigma_{\frac{i}{2}}(e_m)^*$ is a left and right multiplier of $\delta^{\frac{1}{2}}$ and that $\sigma_{\frac{i}{2}}(e_m)^* \delta^{\frac{1}{2}} = \delta^{\frac{1}{2}} \sigma_{\frac{i}{2}}(e_m)^*$.

This implies easily that $\pi(\sigma_{\frac{i}{2}}(e_m))^* \pi(\delta^{\frac{1}{2}}) \subseteq \pi(\delta^{\frac{1}{2}}) \pi(\sigma_{\frac{i}{2}}(e_m))^* \in B(H)$.

So we get that

$$(J\pi(\sigma_{\frac{i}{2}}(e_m))^* J) (J\pi(\delta^{\frac{1}{2}})J) \subseteq (J\pi(\delta^{\frac{1}{2}})J) (J\pi(\sigma_{\frac{i}{2}}(e_m))^* J) \in B(H) .$$

Because $(J\pi(\sigma_{\frac{i}{2}}(e_n))^* J)_{n=1}^\infty$ is a bounded sequence which converges strongly to 1 and $\Lambda(A)$ is dense in H , it is not too difficult to infer from this last equality that the set

$$\langle J\pi(\sigma_{\frac{i}{2}}(e_m))^* J\Lambda(a) \mid n \in \mathbb{N}, a \in \mathcal{N}_\varphi \rangle$$

is a core for $J\pi(\delta^{\frac{1}{2}})J$.

By lemma 8.23 and proposition 6.10, we have for every $a \in \mathcal{N}_\varphi$ and $n \in \mathbb{N}$ that ae_n belongs to $\mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta}$ and $\Lambda(ae_m) = J\pi(\sigma_{\frac{i}{2}}(e_m))^* J\Lambda(a)$. So we see that $\Lambda(\mathcal{N}_\varphi \cap \mathcal{N}_{\varphi_\delta})$ is a core for $J\pi(\delta^{\frac{1}{2}})J$. ■

In a last proposition, we will prove that the strictly positive element δ is uniquely determined in the GNS-representation of φ by the weight φ_δ .

Proposition 8.41 Consider a C^* -algebra A and a KMS-weight φ on A with modular group σ and GNS-construction (H, Λ, π) . Let δ_1, δ_2 be strictly positive elements affiliated with A such that there exists strictly positive numbers λ_1, λ_2 satisfying $\sigma_t(\delta_1) = \lambda_1^t \delta_1$ and $\sigma_t(\delta_2) = \lambda_2^t \delta_2$ for $t \in \mathbb{R}$. If $\varphi_{\delta_1} = \varphi_{\delta_2}$, then $\pi(\delta_1) = \pi(\delta_2)$.

Proof : Because $\varphi_{\delta_1} = \varphi_{\delta_2}$, we have that $\mathcal{N}_{\varphi_{\delta_1}} = \mathcal{N}_{\varphi_{\delta_2}}$ so $\Lambda(\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi_{\delta_1}}) = \Lambda(\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi_{\delta_2}})$.

By lemma 8.40, this gives a common core $\Lambda(\mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi_{\delta_1}})$ for $J\pi(\delta_1^{\frac{1}{2}})J$ and $J\pi(\delta_2^{\frac{1}{2}})J$.

Using lemma 8.39, we have moreover for every $a \in \mathcal{N}_{\varphi} \cap \mathcal{N}_{\varphi_{\delta_1}}$ that

$$\|J\pi(\delta_1^{\frac{1}{2}})J\Lambda(a)\|^2 = \varphi_{\delta_1}(a^*a) = \varphi_{\delta_2}(a^*a) = \|J\pi(\delta_2^{\frac{1}{2}})J\Lambda(a)\|^2.$$

This implies easily that $D(J\pi(\delta_1^{\frac{1}{2}})J) = D(J\pi(\delta_2^{\frac{1}{2}})J)$ and $\|J\pi(\delta_1^{\frac{1}{2}})Jv\| = \|J\pi(\delta_2^{\frac{1}{2}})Jv\|$ for every $v \in D(J\pi(\delta_1^{\frac{1}{2}})J)$.

As a consequence, we get that $J\pi(\delta_1)J = J\pi(\delta_2)J$ which gives us that $\pi(\delta_1) = \pi(\delta_2)$. ■

Corollary 8.42 Consider a C^* -algebra A and a faithful KMS-weight φ on A with modular group σ . Let δ_1, δ_2 be strictly positive elements affiliated with A such that there exists strictly positive numbers λ_1, λ_2 satisfying $\sigma_t(\delta_1) = \lambda_1^t \delta_1$ and $\sigma_t(\delta_2) = \lambda_2^t \delta_2$ for $t \in \mathbb{R}$. If $\varphi_{\delta_1} = \varphi_{\delta_2}$, then $\delta_1 = \delta_2$.

References

- [1] S. BAAJ, Prolongement d'un poids. *C.R. Acad. Sci.* **288** (1979), 1013–1015.
- [2] R.C. BUSBY, Double centralizers and extensions of C^* -algebras. *Trans. Amer. Math. Soc.* **132** (1968), 79–99.
- [3] F. COMBES, Poids sur une C^* -algèbre. *J. Math. pures et appl.* **47** (1968), 57–100.
- [4] F. COMBES, Poids associé à une algèbre hilbertienne à gauche. *Compos. Math.* **23** (1971), 49–77.
- [5] M. ENOCK & J.-M. VALLIN, C^* -algèbres de Kac et algèbres de Kac. *Proc. London Math. Soc.* (3) **66** (1993), 619–650.
- [6] U. HAAGERUP, A density theorem for left Hilbert algebras. Unpublished note.
- [7] U. HAAGERUP, Normal weights on W^* -algebras. *J. Funct. Anal.* **19** (1975), 302–317.
- [8] J. KUSTERMANS & A. VAN DAELE, C^* -algebraic quantum groups arising from algebraic quantum groups. (1996) To appear in *International Journal of Mathematics*.
- [9] J. KUSTERMANS, One-parameter representations on C^* -algebras. In preparation.
- [10] J. KUSTERMANS, Regular C^* -valued weights on C^* -algebras. *Preprint K.U. Leuven* (1997).
- [11] J. KUSTERMANS, Universal C^* -algebraic quantum groups arising from algebraic quantum groups. *Preprint Odense Universitet.* (1997)
- [12] C. LANCE, Hilbert C^* -modules, a toolkit for operator algebraists. Leeds. (1993).
- [13] G.K. PEDERSEN, C^* -algebras and their automorphism groups. Academic Press, London, 1979.

- [14] G.K. PEDERSEN & M. TAKESAKI, The Radon-Nikodym theorem for von Neumann algebras. *Acta Math.* **130** (1973), 53–87.
- [15] J. QUAEGEBEUR & J. VERDING, A construction for weights on C^* -algebras. Dual weights for C^* -crossed products. *Preprint K.U. Leuven* (1994).
- [16] J. QUAEGEBEUR & J. VERDING, Left invariant weights and the left regular corepresentation for locally compact quantum semi-groups. *Preprint K.U. Leuven* (1994).
- [17] S. SAKAI, C^* -algebras and W^* -algebras. Springer-Verlag, Berlin, 1971.
- [18] S. STRATILA & L. ZSIDÓ, Lectures on von Neumann algebras. *Abacus Press, Tunbridge Wells, England* (1979).
- [19] D.C. TAYLOR, The Strict Topology for Double Centralizer Algebras. *Trans. Am. Math. Soc***150** (1970), 633 – 643
- [20] M. TAKESAKI, Theory of Operator Algebras I. *Springer-Verlag, New York* (1979).
- [21] A. VAN DAELE, An Algebraic Framework for Group Duality. (1996) To appear in *Advances of Mathematics*.
- [22] J. VERDING, Weights on C^* -algebras. *Phd-thesis. K.U. Leuven* (1995)
- [23] S.L. WORONOWICZ, From multiplicative unitaries to quantum groups. *Preprint Warszawa* (1995).
- [24] S.L. WORONOWICZ, Unbounded elements affiliated with C^* -algebras and non-compact quantum groups. *Commun. Math. Phys.* **136** (1991), 399–432.
- [25] I. CIORĂNESCU AND L. ZSIDÓ, Analytic generators for one-parameter groups. *Tôhoku Math. Journal* **28** (1976), 327–362.